

Multilayer and Multilevel Cosmological Models based on Solutions of the extended Einstein field equations

Mikhail Batanov-Gaukhman

Ph.D., Associate Professor, Institute No. 2 "Aircraft, rocket engines and power plants",
Federal State Budgetary Educational Institution of Higher Education "Moscow Aviation Institute (National Research University)",
Volokolamsk highway 4, Moscow, Russian Federation (alsignat@yandex.ru)
(Dated: January 27, 2023)

Abstract: This article proposes the multilayer and multilevel cosmological model, on average, balanced with respect to the Einsteinian vacuum. This hierarchical model is based on 10 possible Kottler solutions of the extended Einstein's field equations.

The extension of the Einstein field equations is associated with the addition of an infinite number of Λ_j -terms to these equations with the condition that the series (i.e., the total sum) of these terms converges to zero.

A simplified ten-layer and ten-level case is considered, in which ten spherical formations are sequentially nested into each other like "nesting dolls", while the radii of these spherical formations correspond to the characteristic sizes of a discrete sequence of observed spherical objects: metagalaxy core, galactic cores, stellar (planet) core, biological cells and nuclei of atoms and elementary particles, etc.

As an example, from the general ten-level (hierarchical) solution of extended Einstein's field equations, two antipodal spherical formations with a core radius commensurate with the classical electron radius are singled out. Therefore, the metric-dynamic models of these formations obtained in this way are called «electron» and «positron».

The article is aimed at the development of differential geometry and the program for the complete geometrization of physics by Clifford-Einstein-Wheeler.

Keywords: cosmological model, multilayer and multilevel cosmological model, Kottler metric, Einstein's field equations, metric signature, geometrized physics.

Foreword

This article proposes a multilayer and multilevel cosmological model, i.e. metric-dynamic model of the surrounding world. This model is based on the religio-philosophical "Principles" which the author has drawn largely from Judaism. At the same time, responses to these principles can be found in other religious and philosophical traditions, such as: Hinduism, Taoism, Neoplatonism, Thomism, etc.

Below are the main religious-philosophical Statements that formed the basis of the worldview of the author (the main aspects of the religious-scientific approach to research are set out in the websites <http://alsigna.ru/> , <http://metraphysics.ru/>).

1]. Statement of the Abrahamic Religious Philosophy:

"The ALLHIGHEST Created the Universe from Infinite Nothing"

We do not know, and most likely will never know, what "Infinite Nothing" is. But for a physical and mathematical model, it is enough to assume that this is the "Void" (Vacuum) with unlimited extent and infinite depth. From this Statement, the scientific principle of "Strict Vacuum Balance" originates, from which it follows that everything that arises from the "Emptiness" appears in a mutually opposite form (convexity - concavity, wave - antiwave, particle - antiparticle, Good - Evil, Love - Hatred, etc.), and with full averaging, all Creation is nullified; Within the framework of this concept (corresponding to the program of the complete geometrization of physics by Clifford - Einstein - Wheeler) there is nothing

in the world except: curvature - anti-curvature, rotation - anti-rotation, deformation-anti-deformation and displacement - anti-displacement of space (i.e., Void or Vacuum).

The complete geometrization of physics is the complete exclusion of ideas about matter (or primary matter). The world is a stable Illusion. But the stability of this Grand Illusion is based not on arbitrariness, but on self-closedness, self-consistency, ultimate rationality and optimality; The concept of "mass" and the dimension of the kilogram must be excluded from science.

2]. Statement of the Abrahamic Religious Philosophy:

"The essence of Being is incomprehensible"

From this Statement, we learn that in principle we cannot comprehend the essence of the extension of the world, that is, the substantiality of space (Emptiness, Vacuum) is imperceptible and incomprehensible for us (*Terra incognita*). But what lends itself to our perception is various local and global deformations, curvature, twisting, displacement and other curvature of space (emptiness). Therefore, the true ways of describing the surrounding world can only be geometric (more precisely, space-metric, and more precisely, metric-dynamic, since the curvature of space is constantly changing and moving). We also know for sure that the deformations of the void do not depend on which side or with the help of which (resting or moving) frame of reference we observe them. Therefore, our knowledge is based on the 4-dimensional general covariant differential geometry of the General Theory of Relativity. We also know for sure that any deformations of the void are necessarily accompanied by similar anti-deformations, so that in total or on average they completely compensate for each other's manifestations. For example, local convexity in space cannot arise without its analogous concavity. But if they are separated from each other, or if they have different stable configurations, then they can coexist without annihilating, but on average they should still disappear (i.e., go wrong).

3]. Statement of the Abrahamic Religious Philosophy:

"The ALMIGHTY Creates in the best possible way and Destroys in the worst possible way"

From this Statement originate the two scientific principles of "Least Action" and "Extreme Entropy", which are combined into one principle of "Extreme Average Efficiency". Whence it follows that the dichotomy "Order and House" is balanced. And all the basic physical formulas should be derived on the basis of the probabilistic-variational calculus. This article is based only on the principle of "least action", on the basis of which David Hilbert received the equations of the General theory of relativity of Albert Einstein. But in another article by the author, both extremal principles were applied in the derivation of the Schrödinger equation.

4]. Statement of the Abrahamic Religious Philosophy:

"Everything was created by the ALMIGHTY on the basis of Algorithms for revealing the 4-letter HIS Name יהוה-י (Yud-Key-Vav-Key)"

Rav Gavriel Davidov (by GOD's Permission), during the 2 years of study of the author of this article, reported on the following Algorithms for Revealing the Name יהוה-י [18]: "Have mercy" (Filling), "Rebuy" (Disclosure), "Tree of Ten Sefirot", "Alef-Bet" (Alphabet), Tanta, and a number of Rules for Disclosing Information from the Torah: "Permutation", "Transliteration", "Gematria", "step-by-step coding", etc.

Almost all the main works of the author are built on these Algorithms, including: the present article "Multilayer and multilevel cosmological model of the Universe", "Algebra of signatures", "Light-

geometry of vacuum", "Fully geometrized massless physics", developed metric-dynamical models of all elementary particles included in the Standard Model (except for the Higgs boson), etc.

5]. Statement of the Abrahamic Religious Philosophy:

"EIN SOF, BareHU (ENDLESS) can only give rise to Infinite Creation"

But the Creation is always limited, therefore from this Statement we teach that the INFINITY of the Creative BEGINNING is a continuous (continuous) and limitless INFINITY, and the Infinity of the Created is a discrete, closed (i.e. limited) infinity. From here follows the disk infinity and closedness of the proposed cosmological model.

6]. Statement of the Abrahamic Religious Philosophy:

"The ALLHIGHEST Created Man (i.e., the Universe, the essence of the Tree of Ten Sefirot) in HIS Image", i.e., "Lower is like TOP"

It follows from this Statement that our World is similar to the Higher Spiritual worlds. That is, it is structured according to the principle of the Tree of ten Sefirot. By studying the Structure of the Higher Worlds with the help of Kabbalah, we come to know the structure of our world. And vice versa, comprehending this world in a scientific-empirical way, we cognize the Upper (Spiritual) Worlds.

The main Religious-Philosophical Principles outlined above are the "Guiding Thread" (like GOD's Providence) in the formation of the cosmological model presented below.

In a broader sense, the scientific-religious approach proposed by Algebra of signatures (<http://alsigna.ru/>), is that in those sections of science that are not available for empirical research at the current level of development of measuring technology, as a priori (i.e., the first working) hypothesis, which is the basis of the mathematical model, take the Statements of Religious Sources:

TORAH (תּוֹרָה),	Talmud (תּלְמוּד),
Gospels (Εὐαγγέλιον),	Qur'an (الْقُرْآن),
Avesta (اوستا)	Dao Te Ching (道德經),
Ved (वेद),	Upanishad (उपनिषद्)
etc.	

Further, in the process of deepening and refining the a priori mathematical model based on Religious sources, various aspects that can be empirically verified should be identified. In conclusion, new empirical data should lead to a refinement of the mathematical model and a more specific and in-depth understanding of the Religious Texts.

At the same time, the Religious-scientific approach is a synthesis of the Greek philosophical culture and the Deep Knowledge of the World Religions.

The article below is one of the steps towards Knowledge Synthesis.

7]. Statement of the Abrahamic Religious Philosophy:

"The ALMIGHTY Created Everything from Nothing, but Everything remained Nothing"

The world is there, but on average it is not. That is, convexity and concavity, from which the world is woven, completely compensate each other. The Law of Conservation of the Void.

Numeral - cifra from Arabic صفر (ṣifr) "empty, zero"

1 Background and introduction

ΑΓΓΕΩΜΕΤΡΗΤΟΣ ΜΗΔΕΙΣ ΕΙΣΙΤΩ
 "Let no one ignorant of geometry enter"
 ascribed to Plato (apocryphal)

In 1915, the efforts of Albert Einstein, with the assistance of Marcel Grossmann, and David Hilbert culminated in the derivation of the equations of general relativity [1]

$$R_{ik} - \frac{1}{2}Rg_{ik} = \frac{8\pi G}{c^2}T_{ik}, \quad (1)$$

where g_{ik} are the components of the metric tensor; T_{ik} are the components of the matter energy-momentum tensor; c is the speed of light in vacuum; G is Newton's gravitational constant;

$R = g^{ik}R_{ik}$ is the scalar curvature;

$$R_{ik} = \frac{\partial \Gamma_{ik}^l}{\partial x^l} - \frac{\partial \Gamma_{il}^l}{\partial x^k} + \Gamma_{ik}^l \Gamma_{lm}^m - \Gamma_{il}^m \Gamma_{mk}^l \text{ is Ricci tensor;} \quad (1a)$$

$$\Gamma_{ik}^\lambda = \frac{1}{2}g^{\lambda\mu} \left(\frac{\partial g_{\mu k}}{\partial x^i} + \frac{\partial g_{i\mu}}{\partial x^k} - \frac{\partial g_{ik}}{\partial x^\mu} \right) \text{ is Christoffel symbols.} \quad (1b)$$

However, equations (1) did not allow the possibility of describing the static universe. Therefore, Einstein in 1917, used the property of covariant derivatives:

$$\nabla_j(R_{ik} - \frac{1}{2}Rg_{ik}) = 0, \quad \nabla_j T_{ik} = 0, \quad \nabla_j g_{ik} = 0, \quad (2)$$

and as a result, in the article [2], he wrote down the Einstein's field equations (13a), which is transformed into the formula

$$R_{ik} - \frac{1}{2}Rg_{ik} + \Lambda g_{ik} = \frac{8\pi G}{c^2}T_{ik}, \quad (3)$$

where Λ is a constant, called the "cosmological constant".

In various cosmological models, for example [4,5,6,8, 10,11], it is often assumed

$$\Lambda = 3/r_a^2 \text{ or } \Lambda = -3/r_a^2 \quad (4)$$


where r_a is the radius of some sphere (in particular, the radius of the observable universe); For $T_{ik} = 0$, the case $\Lambda > 0$ corresponds to the de Sitter model, $\Lambda < 0$ corresponds to the anti-de Sitter model.

This article assumes the complete absence of matter ($T_{ik} = 0$), i.e. the vacuum version of the Einstein field equations (3) is considered

$$R_{ik} - \frac{1}{2}Rg_{ik} + \Lambda g_{ik} = 0. \quad (5)$$

Combining Eq.s (5) with the contravariant components of the metric tensor g^{ik} , we obtain

$$g^{ik} \left\{ R_{ik} - \frac{1}{2}Rg_{ik} + \Lambda g_{ik} \right\} = R - \frac{n}{2}R + n\Lambda = 0, \quad (6)$$



where $n = g^{ik}g_{ik}$ is the number of space dimensions.

From Ex. (6) it follows

$$R = \frac{2n}{n-2}\Lambda, \quad (7)$$

in this case, Eq.s (5) takes the form

$$R_{ik} - \frac{n}{n-2}\Lambda g_{ik} + \Lambda g_{ik} = R_{ik} - \frac{2}{n-2}\Lambda g_{ik} = 0.$$

For a 4-dimensional space: $n = 4$, $R = 4\Lambda$, and Eq.s (5) takes the simplest form [4]

$$R_{ik} - \Lambda g_{ik} = 0 \quad (8)$$

$$\text{or} \quad R_{ik} = \Lambda g_{ik} \quad (8a)$$

This equation, taking into account Ex.s (4), can be represented as a system

$$R_{ik} = \pm \frac{3}{r_a^2} g_{ik} = \begin{cases} R_{ik} = \frac{3}{r_a^2} g_{ik}; \\ R_{ik} = -\frac{3}{r_a^2} g_{ik}. \end{cases} \quad (9a)$$

$$(9b)$$

where r_a can also take both positive and negative values ($\pm r_a$).

Eq.s (8) are essentially conservation laws, since the condition (3b.A1) in Appendix 1 is satisfied, and the solutions of these equations describe the metric-dynamic state of stable deformations of local or global areas of vacuum.

The solution of Eq.s (9a) is written in the form of the Kottler metric, which is also called the de Sitter-Schwarzschild solution [4,5,6,12]

$$ds_{Kottler}^2 = \left(1 - \frac{r_b}{r} - \frac{r^2}{r_a^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_b}{r} - \frac{r^2}{r_a^2}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (10)$$

where r_a and r_b are constant parameters of the metric with distance dimension.

In the case: $r_a = \infty$ and $r_b \neq 0$, the Kottler metric (10) turns into the Schwarzschild metric

$$ds_{Schwarzschild}^2 = \left(1 - \frac{r_b}{r}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_b}{r}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2). \quad (11)$$

In another case: $r_a \neq \infty$ and $r_b = 0$, the Kottler metric (10) becomes the de Sitter metric

$$ds_{de\ Sitter}^2 = \left(1 - \frac{r^2}{r_a^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r^2}{r_a^2}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2). \quad (12)$$

In the third case: $r_a = \infty$ and $r_b = 0$, the metric (10) takes the form of the Minkowski metric

$$ds_{Minkowski}^2 = c^2 dt^2 - dr^2 - r^2(d\theta^2 + \sin^2 \theta d\phi^2). \quad (13)$$

28. Die Massenschale.
Wir schließen dieses Kapitel mit einer Betrachtung der Massenschale. Es sei nämlich die homogene Massendichte $\mu = \frac{\rho}{c^2}$ nicht mehr über die Vollkugel $0 \leq r \leq a$ ausgebreitet, sondern nur über die Schale, $0 < b \leq r \leq a$. Der einzige Unterschied gegen früher ist, daß jetzt in der Materie $\frac{1}{g_1} = 1 - \frac{r^2}{R^2} + \frac{\text{const}}{r}$ $b \leq r \leq a$ genommen werden kann. Für den Innenraum $0 \leq r \leq b$ haben wir $\frac{1}{g_1} = 1 + \frac{\text{const}}{r}$ und hier muß wieder die Konstante verschwinden; also $g_1 = 1$ für $0 \leq r \leq b$. Andererseits ist $\frac{1}{g_1} = 1 - \frac{2km}{c^2} \cdot \frac{1}{r}$ für $a \leq r < \infty$, da aber jetzt $\frac{2km}{c^2} = \frac{8\pi k}{3c^2} \epsilon (a^3 - b^3) = \frac{8\pi k}{3} (a^3 - b^3)$, folgt bei stetigem Anschluß der g_1 -Werte für $r = b$ bzw. $r = a$ wieder $ks = \frac{3}{R^2}$, womit die analogen Schlußfolgerungen wie vorhin auftreten. Die Lösung ist jetzt, wie man leicht findet: $g_1 = 1, \quad g_4 = c^2$ für $0 \leq r \leq b$, $\frac{1}{g_1} = 1 - \frac{r^2}{R^2} + \frac{b^2}{R^2} \cdot \frac{1}{r}, \quad g_4 = c^2 \left(1 - \frac{r^2}{R^2} + \frac{b^2}{R^2} \cdot \frac{1}{r}\right)$ für $b \leq r \leq a$, $\frac{1}{g_1} = 1 - \frac{a^2 - b^2}{R^2} \cdot \frac{1}{r}, \quad g_4 = c^2 \left(1 - \frac{a^2 - b^2}{R^2} \cdot \frac{1}{r}\right)$ für $a \leq r < \infty$. Wir haben also das wichtige Resultat:

2 Generalized Kottler metrics

In fact, Friedrich Kottler wrote down in [3] not the metric

(10), but the metric of the form

$$ds_{\text{Kot}}^2 = - \left(1 + \frac{r_b}{r} - \frac{r^2}{r_a^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_b}{r} - \frac{r^2}{r_a^2} \right)} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2)$$

this can be easily seen on page 443 of that article.

Article [3] was published in March 1918, that is, almost immediately after the publication of Einstein's GR, so the metrics given below will be called the generalized Kottler metrics.

Since the metric (10) is used in many cosmological models, for example, [4,5,6,7,8,9,10,11,12], in this article it is proposed to

pay attention to the fact that the parameters $3/r_a^2$ and r_b can take both positive and negative values ($\pm 3/r_a^2$ and $\pm r_b$), so the system of Eq.s (9) generally has the following 10 mutually irreducible solutions:

- with signature (+ - - -)

$$ds_1^{(-)2} = \left(1 - \frac{r_{b1}}{r} + \frac{r^2}{r_{a1}^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_{b1}}{r} + \frac{r^2}{r_{a1}^2} \right)} - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (14)$$

$$ds_2^{(-)2} = \left(1 + \frac{r_{b2}}{r} - \frac{r^2}{r_{a2}^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_{b2}}{r} - \frac{r^2}{r_{a2}^2} \right)} - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (15)$$

$$ds_3^{(-)2} = \left(1 - \frac{r_{b3}}{r} - \frac{r^2}{r_{a3}^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_{b3}}{r} - \frac{r^2}{r_{a3}^2} \right)} - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (18)$$

$$ds_4^{(-)2} = \left(1 + \frac{r_{b4}}{r} + \frac{r^2}{r_{a4}^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_{b4}}{r} + \frac{r^2}{r_{a4}^2} \right)} - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (19)$$

$$ds_5^{(-)2} = c^2 dt^2 - dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2); \quad (20)$$

- and with signature (- + + +)

$$ds_1^{(+)2} = - \left(1 - \frac{r_{b1}}{r} + \frac{r^2}{r_{a1}^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_{b1}}{r} + \frac{r^2}{r_{a1}^2} \right)} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (21)$$

$$ds_2^{(+)2} = - \left(1 + \frac{r_{b2}}{r} - \frac{r^2}{r_{a2}^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_{b2}}{r} - \frac{r^2}{r_{a2}^2} \right)} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (22)$$

$$ds_3^{(+)2} = - \left(1 - \frac{r_{b3}}{r} - \frac{r^2}{r_{a3}^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_{b3}}{r} - \frac{r^2}{r_{a3}^2} \right)} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (23)$$

$$ds_4^{(+)2} = - \left(1 + \frac{r_{b4}}{r} + \frac{r^2}{r_{a4}^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_{b4}}{r} + \frac{r^2}{r_{a4}^2} \right)} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (24)$$

$$ds_5^{(+)2} = - c^2 dt^2 + dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (25)$$

where $r_{a1}, r_{a2}, r_{a3}, r_{a4}$ and $r_{b1}, r_{b2}, r_{b3}, r_{b4}$ are parameters of the cosmological model.

All ten metrics (14) – (25) are solutions of one system of Eq.s (9), which determines the metric-dynamic state of the same region of space. Therefore, we can assume that these solutions describe the metric-dynamic state of the ten "layers" of this area. In this case, the additive overlay (i.e., sum or average) of all ten metrics

$$\sum_{i=1}^5 s_i^{(-)2} + \sum_{j=1}^5 s_j^{(+)2} = 0 \quad (26)$$

leads to two more trivial solutions of Eq.s (9) with mutually opposite signatures (+ ---) and (- +++)

$$0 = 0 \cdot c^2 dt^2 - 0 \cdot dr^2 - 0 \cdot r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (27)$$

$$0 = -0 \cdot c^2 dt^2 + 0 \cdot dr^2 + 0 \cdot r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (28)$$

This can be easily seen if we substitute the zero components of the metric tensor $g_{ik} = 0$ from metrics (27) and (28) into Eq.s (9) taking into account Ex.s (1a) and (1b).

All 12 solutions of the system Einstein's field equations (9) will be taken into account in the multi-layer cosmological model.

Before discussing the possibility of additive superposition (i.e., summation or averaging) of metric layers on each other, we note the following important circumstance.

Addition of two quadratic forms

$$ds_a^2 + ds_b^2 = ds_{ab}^2, \quad (29)$$

or their averaging $1/2 ds_a^2 + 1/2 ds_b^2 = 1/2 ds_{ab}^2 \quad (29a)$

resembles the Pythagorean theorem (see Figure 1a) $a^2 + b^2 = c^2$.

This means that the segments ds_a and ds_b of the corresponding lines s_a and s_b are always mutually perpendicular to each other ($ds_a \perp ds_b$). This is possible only if the lines s_a and s_b form a double helix (see Figure 1b), which can be described by a system of two complex conjugate numbers

$$ds_{ab} = ds_a + i ds_b, \quad (30)$$

$$ds_{ab}^* = ds_a' - i ds_b'. \quad (30a)$$

The product of complex numbers (30) and (30a), under the condition $ds_a = ds_a'$ и $s_b = ds_b'$, is equal to the quadratic form (29).

Such a spiral will be called a 4-braid (i.e., a bundle consisting of 4 interlaced lines s_a, s_b, s_a', s_b'), wherein:

- the line s_a belongs to the outer side of the a -th affine space;
- the line s_a' belongs to the inner side of the a -th affine space;
- the line s_b belongs to the outside of the b -th affine space;
- the line s_b' belongs to the inner side of the b -th affine space.




Fig. 1. Double helix of lines s_a and s_b



Fig. 2. Illustration of a 4 line s_a, s_b, s_a', s_b' spiral

The addition of quadratic forms (29) or their averaging (29a) depends on the model representation. If we assume that two metric spaces are additively superimposed on each other, then the addition operation (29) should be used. If we assume that a metric space can be in a state with the metric s_a^2 or with the metric s_b^2 with equal probability, then this corresponds to the quantum mechanical approach, and the averaging operation (29a) should be used.

At this stage of the study, it is difficult to decide which operation "addition" (29) or "averaging" (29a) should be used. However, the quantum mechanical approach provides additional opportunities for probability theory and does not require an answer to the question: "Is the sum of solutions (for example, s_a^2 and s_b^2) of Einstein's non-linear differential equations also a solution of this equations?". Therefore, the quantum mechanical approach looks much more preferable, especially when the sum of solutions is also a solution to a nonlinear equation. Arguments in favor of the second approach are presented in Appendix 1, see Ex.s (4.A1) – (32.A1).

Now, for example, consider the averaging of four metrics (14) – (20)

$$ds_{1-4}^{(-)2} = \frac{1}{4} (ds_1^{(-)2} + ds_2^{(-)2} + ds_3^{(-)2} + ds_4^{(-)2}). \quad (31)$$

As a result, we obtain the averaged metric

$$ds_{1-4}^{(-)2} = f(r)c^2 dt^2 - k(r)dr^2 - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (32)$$

$$\text{where } f(r) = \frac{1}{4} \left[\left(1 - \frac{r_{b1}}{r} + \frac{r^2}{r_{a1}^2} \right) + \left(1 + \frac{r_{b2}}{r} - \frac{r^2}{r_{a2}^2} \right) + \left(1 - \frac{r_{b3}}{r} - \frac{r^2}{r_{a3}^2} \right) + \left(1 + \frac{r_{b4}}{r} + \frac{r^2}{r_{a4}^2} \right) \right], \quad (33)$$

$$k(r) = \frac{1}{4} \left[\frac{1}{\left(1 - \frac{r_{b1}}{r} + \frac{r^2}{r_{a1}^2} \right)} + \frac{1}{\left(1 + \frac{r_{b2}}{r} - \frac{r^2}{r_{a2}^2} \right)} + \frac{1}{\left(1 - \frac{r_{b3}}{r} - \frac{r^2}{r_{a3}^2} \right)} + \frac{1}{\left(1 + \frac{r_{b4}}{r} + \frac{r^2}{r_{a4}^2} \right)} \right]. \quad (34)$$

By analogy with (29) – (30), such a metric space is based on the interweaving of four affine extensions, which respectively belong to 8 linear forms $ds_i^{(-)}$ and $ds_i^{(-)'}$, i.e. segments of 8 lines twisted into an 8-braid.

Such an 8-braid is described by a system of two complex conjugate quaternions:

$$ds_{1-4}^{(-)} = \frac{1}{\sqrt{4}} \left(ds_1^{(-)} + i ds_2^{(-)} + j ds_3^{(-)} + k ds_4^{(-)} \right), \quad (35)$$

$$ds_{1-4}^{(-)*} = \frac{1}{\sqrt{4}} \left(ds_1^{(-)'} - i ds_2^{(-)'} - j ds_3^{(-)'} - k ds_4^{(-)'} \right), \quad (35a)$$

whose product is equal to the metric (32).

On Figure 3 shows an illustration of the interweaving of several affine subspaces that form a multilayer metric space.

Properties of intertwined affine subspaces and multilayer metric spaces with signatures $(+ - - -)$ and $(- + + +)$ corresponding to the "vacuum balance" condition




Fig. 3. Illustration of the interweaving of several affine subspaces forming a multi-layer metric space

$$(+---) + (-+++)=0 \quad (36)$$

detailed in the "Algebra of signatures" [16,17].

In the same papers [16,17], a more detailed analysis was carried out, considering metric spaces with all sixteen possible signatures:

$$\begin{array}{cccc} (+ + + +) & (+ + + -) & (- + + -) & (+ + - +) \\ (- - - +) & (- + + +) & (- - + +) & (- + - +) \\ (+ - - +) & (+ + - -) & (+ - - -) & (+ - + +) \\ (- - + -) & (+ - + -) & (- + - -) & (- - - -), \end{array} \quad (37)$$

which in total correspond to the principle of "full vacuum (i.e., zero) balance",

$$\begin{array}{rclcl} 0 = & \underline{(0 \ 0 \ 0 \ 0)} & + & \underline{(0 \ 0 \ 0 \ 0)} & = 0 \\ 0 = & (+ \ + \ + \ +) & + & (- \ - \ - \ -) & = 0 \\ 0 = & (- \ - \ - \ +) & + & (+ \ + \ + \ -) & = 0 \\ 0 = & (+ \ - \ - \ +) & + & (- \ + \ + \ -) & = 0 \\ 0 = & (- \ - \ + \ -) & + & (+ \ + \ - \ +) & = 0 \\ 0 = & (+ \ + \ - \ -) & + & (- \ - \ + \ +) & = 0 \\ 0 = & (- \ + \ - \ -) & + & (+ \ - \ + \ +) & = 0 \\ 0 = & (+ \ - \ + \ -) & + & (- \ + \ - \ +) & = 0 \\ 0 = & \underline{(- \ + \ + \ +)} & + & \underline{(+ \ - \ - \ -)} & = 0 \\ 0 = & (0 \ 0 \ 0 \ 0)_+ & + & (0 \ 0 \ 0 \ 0)_+ & = 0 \end{array} \quad (37a)$$

In this expression, called in [16,17] ranking, the summation of the signs "+" and "-" is performed both in columns and in rows.

The simplified "vacuum balance" (36) follows from the ranking Ex. (37a) [16,17]

$$\begin{array}{rclcl} (+ \ + \ + \ +) & + & (- \ - \ - \ -) & = 0 \\ (- \ - \ - \ +) & + & (+ \ + \ + \ -) & = 0 \\ (+ \ - \ - \ +) & + & (- \ + \ + \ -) & = 0 \\ (- \ - \ + \ -) & + & (+ \ + \ - \ +) & = 0 \\ (+ \ + \ - \ -) & + & (- \ - \ + \ +) & = 0 \\ (- \ + \ - \ -) & + & (+ \ - \ + \ +) & = 0 \\ \underline{(+ \ - \ + \ -)} & + & \underline{(- \ + \ - \ +)} & = 0 \\ (+ \ - \ - \ -)_+ & + & (- \ + \ + \ +)_+ & = 0. \end{array} \quad (37b)$$

That is, the additive superposition (in this case, the addition of signs of signatures by columns) of seven metric spaces with signatures in the denominator of the left column of the ranking Ex. (37b), form a Minkowski space with signature (+---).

Whereas the additive superposition of seven metric spaces with signatures in the denominator of the right column of the ranking Ex. (37b), form an anti-Minkowski space with the opposite signature (-+++).

Accounting for all sixteen metric spaces with signatures (37) can significantly enrich the cosmological models of the universe (see Appendix 2 and also [16,17]).

3 Extended Einstein's field equations

In the previous paragraphs of this article, we considered a set of solutions to the Einstein field equations (8), well known to specialists. In this paragraph, for the first time, it is proposed to consider an extended version of this equations.

Recall that Einstein, in order to write Eq.s (3), used the following property of the metric tensor

$$\Lambda \nabla_j g_{ik} = \nabla_j \Lambda g_{ik} = 0, \quad (38)$$

however, it is obvious that the covariant derivative of the infinite series $\Lambda_l \nabla_j g_{ik}$ is also equal to zero

$$\nabla_j (\Lambda_1 g_{ik} + \Lambda_2 g_{ik} + \Lambda_3 g_{ik} + \dots + \Lambda_\infty g_{ik}) = \Lambda_1 \nabla_j g_{ik} + \Lambda_2 \nabla_j g_{ik} + \dots + \Lambda_\infty \nabla_j g_{ik} = 0, \quad (39)$$

where $\Lambda_1, \Lambda_2, \dots, \Lambda_\infty$ are constants that can take both positive ($\Lambda_i > 0$) and negative ($\Lambda_i < 0$) values.

Therefore, we use the same method that Einstein used to introduce the Λ -term into Eq.s (1) [2], and write the equations

$$R_{ik} - \frac{1}{2} R g_{ik} + \Lambda_1 g_{ik} + \Lambda_2 g_{ik} + \Lambda_3 g_{ik} + \dots + \Lambda_\infty g_{ik} = 0,$$

or in a more compact form

$$R_{ik} - \frac{1}{2} R g_{ik} + g_{ik} \sum_{j=1}^{\infty} \Lambda_j = 0, \quad (40)$$

where $\Lambda_j = 3/r_{aj}^2$ or $-3/r_{aj}^2$, here r_{aj} is the radius of the j -th spherical formation.

If the sum of the series $\Lambda_1 + \Lambda_2 + \Lambda_3 + \dots + \Lambda_\infty$ converges to a constant number Λ_0 , i.e., if

$$\sum_{j=1}^{\infty} \Lambda_j = \Lambda_0, \quad (41)$$

then Eq.s (40) takes the form of Eq.s (5)

$$R_{ik} - \frac{1}{2} R g_{ik} + \Lambda_0 g_{ik} = 0, \quad (42)$$

which, like (6) – (7), is reduced to the form similar to Eq.s (8)

$$R_{ik} - g_{ik} \Lambda_0 = 0. \quad (43)$$

Therefore, the solutions of Eq.s (43) practically coincide with the solutions (14) – (25):

- with signature (+ ---)

$$ds_1^{(-)2} = \left(1 - \frac{r_f}{r} + \frac{\Lambda_0 r^2}{3}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_f}{r} + \frac{\Lambda_0 r^2}{3}\right)} - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (44)$$

$$ds_2^{(-)2} = \left(1 + \frac{r_f}{r} - \frac{\Lambda_0 r^2}{3}\right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_f}{r} - \frac{\Lambda_0 r^2}{3}\right)} - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (45)$$

$$ds_3^{(-)2} = \left(1 - \frac{r_f}{r} - \frac{\Lambda_0 r^2}{3}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_f}{r} - \frac{\Lambda_0 r^2}{3}\right)} - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (46)$$

$$ds_4^{(-)2} = \left(1 + \frac{r_f}{r} + \frac{\Lambda_0 r^2}{3}\right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_f}{r} + \frac{\Lambda_0 r^2}{3}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (47)$$

$$ds_5^{(-)2} = c^2 dt^2 - dr^2 - r^2(d\theta^2 + \sin^2 \theta d\phi^2); \quad (48)$$

- and with signature $(-+++)$

$$ds_5^{(+)2} = -c^2 dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (49)$$

$$ds_4^{(+)2} = -\left(1 + \frac{r_f}{r} + \frac{\Lambda_0 r^2}{3}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_f}{r} + \frac{\Lambda_0 r^2}{3}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (50)$$

$$ds_3^{(+)2} = -\left(1 - \frac{r_f}{r} - \frac{\Lambda_0 r^2}{3}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_f}{r} - \frac{\Lambda_0 r^2}{3}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (51)$$

$$ds_2^{(+)2} = -\left(1 + \frac{r_f}{r} - \frac{\Lambda_0 r^2}{3}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_f}{r} - \frac{\Lambda_0 r^2}{3}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (52)$$

$$ds_1^{(+)2} = -\left(1 - \frac{r_f}{r} + \frac{\Lambda_0 r^2}{3}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_f}{r} + \frac{\Lambda_0 r^2}{3}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (53)$$

where Λ_0 and r_f are the zero results of the summation of alternating series

$$\Lambda_0 = \sum_{j=1}^{\infty} \Lambda_j = \sum_{j=1}^{\infty} (-1)^j \frac{3N_j}{r_{aj}^2} = \sum_{n=2j}^{\infty} \frac{3N_n}{r_{an}^2} + \sum_{m=2j-1}^{\infty} \left(-\frac{3N_m}{r_{am}^2}\right) = 0, \quad (54)$$

$$r_f = \sum_{j=1}^{\infty} (-1)^j r_{bj} = \sum_{n=2j}^{\infty} r_{bn} + \sum_{m=2j-1}^{\infty} (-r_{bm}) = 0, \quad (55)$$

where N_j are the dimensionless corrective parameters of the considered cosmological model.

Such a cosmological model is the most optimal, since in the case of $\Lambda_0 = 0$ (54) the extended Einstein field equations as a whole take on the simplest form

$$R_{ik} = 0, \quad (55a)$$

with four Schwarzschild solutions

$$ds_1^{(-)2} = \left(1 - \frac{r_f}{r}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_f}{r}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2),$$

$$ds_2^{(-)2} = \left(1 + \frac{r_f}{r}\right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_f}{r}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2),$$

$$ds_4^{(+)2} = -\left(1 + \frac{r_f}{r}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_f}{r}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2),$$

$$ds_3^{(+)2} = -\left(1 - \frac{r_f}{r}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_f}{r}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2),$$

which, when condition (55) is met, turn into two Minkowski solutions (20) and (25), which is consistent with the principle of “maximum rationality” and the principle of “total average absence” (i.e., observance of the “vacuum balance”).

At the same time, it is possible that there is a slight imbalance between the positive and negative members of the series (54)

$$\Lambda_0 = \sum_{n=2j}^{\infty} \frac{3N_n}{r_{an}^2} + \sum_{m=2j-1}^{\infty} \left(-\frac{3N_m}{r_{am}^2} \right) \neq 0, \quad (55b)$$

$$r_f = \sum_{n=2j}^{\infty} r_{bn} + \sum_{m=2j-1}^{\infty} (-r_{bm}) = 0, \quad (55c)$$

where $j=1,2,3,\dots$

For example, it is possible that the imbalance exists, but is extremely small.

$$\lim_{n,m \rightarrow \infty} \Lambda_0 = \Lambda_{\Sigma} = 1.0905 \cdot 10^{-52} \text{ m}^{-2},$$

i.e., tend to the value published by the Plank collaboration [14] for the Standard cosmological model Lambda-CDM; where Λ_{Σ} is the total cosmological constant (TCC).

Then Eq.s (43) takes the form

$$R_{ik} = \Lambda_{\Sigma} g_{ik}, \quad (55d)$$

with four de Sitter solutions

$$\begin{aligned} ds_1^{(-)2} &= \left(1 + \frac{\Lambda_{\Sigma} r^2}{3}\right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{\Lambda_{\Sigma} r^2}{3}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \\ ds_2^{(-)2} &= \left(1 - \frac{\Lambda_{\Sigma} r^2}{3}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{\Lambda_{\Sigma} r^2}{3}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \\ ds_4^{(+)2} &= -\left(1 + \frac{\Lambda_{\Sigma} r^2}{3}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{\Lambda_{\Sigma} r^2}{3}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \\ ds_3^{(+)2} &= -\left(1 - \frac{\Lambda_{\Sigma} r^2}{3}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{\Lambda_{\Sigma} r^2}{3}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2). \end{aligned}$$

In this case, according to Ex.s (7) and (4) $4\Lambda_{\Sigma} = R$, so $R = \frac{12}{r_{a\Sigma}^2}$, where

$$r_{a\Sigma} = \sqrt{\frac{R}{12}} = \sqrt{\frac{\Lambda_{\Sigma}}{3}} \approx 1,6586 \cdot 10^{26} \text{ m}. \quad (55e)$$

If there is a predominance of positive terms in Ex. (55b), then $\Lambda_{\Sigma} > 0$ (de Sitter universe); if negative terms predominate, then $\Lambda_{\Sigma} < 0$ (anti - de Sitter universe).

At the same time, in the author's opinion, the "vacuum balance" cannot be violated, since only mutually opposite entities can appear from the void (for example, convexity - concavity, wave - antiwave, particle-antiparticle, etc.). Therefore, if an imbalance (55b) of type $\Lambda_{\Sigma} > 0$ exists, then a counter-imbalance of type $\Lambda_{\Sigma} < 0$ must also coexist.

It is possible to restore the averaged "vacuum balance" in such a situation if we assume that the sign of the total cosmological constant (TCC) periodically changes with time. For example, it can be assumed that the value of the TCC fluctuates according to a sinusoidal law with an amplitude $\Lambda_{\Sigma} \approx 10^{-52} \text{ m}^{-2}$. Then, on average, the "vacuum balance" is restored

$$R_{ik} = g_{ik} \overline{\Lambda_{\Sigma} \sin(2\pi f_u t)} = 0, \quad (55f)$$

where f_u is the cosmological oscillation frequency.

4 Closed ten-layer and ten-level cosmological model

In the previous section, it was shown that the extended Einstein field equations can contain an infinite number of $\pm\Lambda_j$ -terms, provided that the sum of all these terms must tend to zero to maintain the “vacuum balance”.

Further, it will be shown that the cosmological model based on the extended Einstein field equations (43), taking into account Ex.s (54) and (55) [or (55b) and (55c)], can be a closed spherical space filled with an infinite number of spherical formations (“bubbles”) and opposite anti-spherical formations (“anti-bubbles”) with different radii r_{aj} , inside which there is an infinite number of smaller “bubbles” and “anti-bubbles” with different radii r_{bj} , and so it continues to infinity from level to level (see Figure 4).

At this point, for simplicity, we single out only ten levels with two mutually opposite spherical formations (i.e., a “bubble” and an “anti-bubble”) at each level.

In other words, to identify the main parameters of the multilevel cosmological model, we study a special case when, instead of infinite series (54) and (55), we use a simplified series limited by ten mutually opposite pairs of terms:

$$\Lambda_{0(10)} = \sum_{j=1}^{10} \Lambda_j = \sum_{j=1}^{10} \frac{3}{r_j^2} + \sum_{j=1}^{10} \left(-\frac{3}{r_j^2}\right) = 0, \quad (56)$$

$$r_{f(10)} = \sum_{k=1}^{10} r_j + \sum_{j=1}^{10} (-r_j) = 0. \quad (57)$$

Consider separately the series with positive and negative terms

$$r_{f(10)} = \sum_{j=1}^{10} r_j, \quad \Lambda_{0(10)} = 3 \sum_{j=1}^{10} \frac{1}{r_j^2}; \quad (58)$$

$$r_{f(-10)} = \sum_{j=1}^{10} (-r_j), \quad \Lambda_{0(-10)} = 3 \sum_{k=1}^{10} \left(-\frac{1}{r_j^2}\right). \quad (59)$$

Let's substitute series (58) into metrics (44) – (48) instead of series (56) and (57) and take into account that we can write:

$$\begin{aligned} 1 - \frac{r_{f(10)}}{r} + \frac{\Lambda_{0(10)} r^2}{3} &= 1 - \frac{r_1 + r_2 + \dots + r_{10}}{r} + \left(\frac{1}{r_1^2} + \frac{1}{r_2^2} + \dots + \frac{1}{r_{10}^2}\right) r^2 = \\ &= \left(1 - \frac{r_{10}}{r} + \frac{r^2}{r_{10}^2}\right) - \left(1 + \frac{r_9}{r} - \frac{r^2}{r_9^2}\right) + \left(1 - \frac{r_8}{r} + \frac{r^2}{r_8^2}\right) - \left(1 + \frac{r_7}{r} - \frac{r^2}{r_7^2}\right) + \left(1 - \frac{r_6}{r} + \frac{r^2}{r_6^2}\right) - \left(1 + \frac{r_5}{r} - \frac{r^2}{r_5^2}\right) + \\ &+ \left(1 - \frac{r_4}{r} + \frac{r^2}{r_4^2}\right) - \left(1 + \frac{r_3}{r} - \frac{r^2}{r_3^2}\right) + \left(1 - \frac{r_2}{r} + \frac{r^2}{r_2^2}\right) - \left(1 + \frac{r_1}{r} - \frac{r^2}{r_1^2}\right) \end{aligned} \quad (60)$$




Fig. 4. Illustration of a sequence of “bubbles” within “bubbles” within “bubbles”...

$$\begin{aligned}
1 + \frac{r_{f(10)}}{r} - \frac{A_{0(10)}r^2}{3} &= 1 + \frac{r_1 + r_2 + \dots + r_{10}}{r} - \left(\frac{1}{r_1^2} + \frac{1}{r_2^2} + \dots + \frac{1}{r_{10}^2} \right) r^2 = \\
&= \left(1 + \frac{r_{10}}{r} - \frac{r^2}{r_9^2} \right) - \left(1 - \frac{r_9}{r} + \frac{r^2}{r_8^2} \right) + \left(1 + \frac{r_8}{r} - \frac{r^2}{r_7^2} \right) - \dots + \left(1 + \frac{r_2}{r} - \frac{r^2}{r_1^2} \right) - \left(1 - \frac{r_1}{r} + \frac{r^2}{r_{10}^2} \right), \quad (61)
\end{aligned}$$

$$\begin{aligned}
1 - \frac{r_{f(10)}}{r} - \frac{A_{0(10)}r^2}{3} &= 1 - \frac{r_1 + r_2 + \dots + r_{10}}{r} - \left(\frac{1}{r_1^2} + \frac{1}{r_2^2} + \dots + \frac{1}{r_{10}^2} \right) r^2 = \\
&= \left(1 - \frac{r_{10}}{r} - \frac{r^2}{r_9^2} \right) - \left(1 + \frac{r_9}{r} + \frac{r^2}{r_8^2} \right) + \left(1 - \frac{r_8}{r} - \frac{r^2}{r_7^2} \right) - \dots + \left(1 - \frac{r_2}{r} - \frac{r^2}{r_1^2} \right) - \left(1 + \frac{r_1}{r} + \frac{r^2}{r_{10}^2} \right), \quad (62)
\end{aligned}$$

$$\begin{aligned}
1 + \frac{rf(10)}{r} + \frac{\Lambda_{0(10)}r^2}{3} &= 1 + \frac{r_1+r_2+\dots+r_{10}}{r} + \left(\frac{1}{r_1^2} + \frac{1}{r_2^2} + \dots + \frac{1}{r_{10}^2}\right)r^2 = \\
&= \left(1 + \frac{r_{10}}{r} + \frac{r^2}{r_9^2}\right) - \left(1 - \frac{r_9}{r} - \frac{r^2}{r_8^2}\right) + \left(1 + \frac{r_8}{r} + \frac{r^2}{r_7^2}\right) - \dots + \left(1 + \frac{r_2}{r} + \frac{r^2}{r_1^2}\right) - \left(1 - \frac{r_1}{r} - \frac{r^2}{r_{10}^2}\right). \quad (63)
\end{aligned}$$

As a result, we obtain five metrics with signature $(+---)$:

$$\begin{aligned}
ds_1^{(-)2} = & \left\{ \left(1 - \frac{r_{10}}{r} + \frac{r^2}{r_9^2}\right) - \left(1 + \frac{r_9}{r} - \frac{r^2}{r_8^2}\right) + \left(1 - \frac{r_8}{r} + \frac{r^2}{r_7^2}\right) - \left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) + \left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) - \right. \\
& \left. - \left(1 + \frac{r_5}{r} - \frac{r^2}{r_4^2}\right) + \left(1 - \frac{r_4}{r} + \frac{r^2}{r_3^2}\right) - \left(1 + \frac{r_3}{r} - \frac{r^2}{r_2^2}\right) + \left(1 - \frac{r_2}{r} + \frac{r^2}{r_1^2}\right) - \left(1 + \frac{r_1}{r} - \frac{r^2}{r_{10}^2}\right) \right\} c^2 dt^2 - \\
& - \left\{ \left(1 - \frac{r_{10}}{r} + \frac{r^2}{r_9^2}\right) - \left(1 + \frac{r_9}{r} - \frac{r^2}{r_8^2}\right) + \left(1 - \frac{r_8}{r} + \frac{r^2}{r_7^2}\right) - \left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) + \left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) - \right. \\
& \left. - \left(1 + \frac{r_5}{r} - \frac{r^2}{r_4^2}\right) + \left(1 - \frac{r_4}{r} + \frac{r^2}{r_3^2}\right) - \left(1 + \frac{r_3}{r} - \frac{r^2}{r_2^2}\right) + \left(1 - \frac{r_2}{r} + \frac{r^2}{r_1^2}\right) - \left(1 + \frac{r_1}{r} - \frac{r^2}{r_{10}^2}\right) \right\}^{-1} dr^2 - \\
& - r^2(d\theta^2 + \sin^2 \theta d\phi^2),
\end{aligned} \tag{64}$$

$$ds_2^{(-)2} = \left\{ \left(1 + \frac{r_{10}}{r} - \frac{r^2}{r_9^2}\right) - \left(1 - \frac{r_9}{r} + \frac{r^2}{r_8^2}\right) + \left(1 + \frac{r_8}{r} - \frac{r^2}{r_7^2}\right) - \left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) + \left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) - \right. \\ \left. - \left(1 - \frac{r_5}{r} + \frac{r^2}{r_4^2}\right) + \left(1 + \frac{r_4}{r} - \frac{r^2}{r_3^2}\right) - \left(1 - \frac{r_3}{r} + \frac{r^2}{r_2^2}\right) + \left(1 + \frac{r_2}{r} - \frac{r^2}{r_1^2}\right) - \left(1 - \frac{r_1}{r} + \frac{r^2}{r_{10}^2}\right) \right\} c^2 dt^2 - \\ - \left\{ \left(1 + \frac{r_{10}}{r} - \frac{r^2}{r_9^2}\right) - \left(1 - \frac{r_9}{r} + \frac{r^2}{r_8^2}\right) + \left(1 + \frac{r_8}{r} - \frac{r^2}{r_7^2}\right) - \left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) + \left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) - \right. \\ \left. - \left(1 - \frac{r_5}{r} + \frac{r^2}{r_4^2}\right) + \left(1 + \frac{r_4}{r} - \frac{r^2}{r_3^2}\right) - \left(1 - \frac{r_3}{r} + \frac{r^2}{r_2^2}\right) + \left(1 + \frac{r_2}{r} - \frac{r^2}{r_1^2}\right) - \left(1 - \frac{r_1}{r} + \frac{r^2}{r_{10}^2}\right) \right\}^{-1} dr^2 - \\ - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (65)$$

$$\begin{aligned}
ds_3^{(-)2} = & \left\{ \left(1 - \frac{r_{10}}{r} - \frac{r^2}{r_9^2}\right) - \left(1 + \frac{r_9}{r} + \frac{r^2}{r_8^2}\right) + \left(1 - \frac{r_8}{r} - \frac{r^2}{r_7^2}\right) - \left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) + \left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) - \right. \\
& \left. - \left(1 + \frac{r_5}{r} + \frac{r^2}{r_4^2}\right) + \left(1 - \frac{r_4}{r} - \frac{r^2}{r_3^2}\right) - \left(1 + \frac{r_3}{r} + \frac{r^2}{r_2^2}\right) + \left(1 - \frac{r_2}{r} - \frac{r^2}{r_1^2}\right) - \left(1 + \frac{r_1}{r} + \frac{r^2}{r_{10}^2}\right) \right\} c^2 dt^2 - \\
& - \left\{ \left(1 - \frac{r_{10}}{r} - \frac{r^2}{r_9^2}\right) - \left(1 + \frac{r_9}{r} + \frac{r^2}{r_8^2}\right) + \left(1 - \frac{r_8}{r} - \frac{r^2}{r_7^2}\right) - \left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) + \left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) - \right. \\
& \left. - \left(1 + \frac{r_5}{r} + \frac{r^2}{r_4^2}\right) + \left(1 - \frac{r_4}{r} - \frac{r^2}{r_3^2}\right) - \left(1 + \frac{r_3}{r} + \frac{r^2}{r_2^2}\right) + \left(1 - \frac{r_2}{r} - \frac{r^2}{r_1^2}\right) - \left(1 + \frac{r_1}{r} + \frac{r^2}{r_{10}^2}\right) \right\}^{-1} dr^2 - \\
& - r^2(d\theta^2 + \sin^2\theta d\phi^2), \tag{66}
\end{aligned}$$

$$\begin{aligned}
ds_4^{(-)2} = & \left\{ \left(1 + \frac{r_{10}}{r} + \frac{r^2}{r_9^2} \right) - \left(1 - \frac{r_9}{r} - \frac{r^2}{r_8^2} \right) + \left(1 + \frac{r_8}{r} + \frac{r^2}{r_7^2} \right) - \left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2} \right) + \left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2} \right) - \right. \\
& \left. - \left(1 - \frac{r_5}{r} - \frac{r^2}{r_4^2} \right) + \left(1 + \frac{r_4}{r} + \frac{r^2}{r_3^2} \right) - \left(1 - \frac{r_3}{r} - \frac{r^2}{r_2^2} \right) + \left(1 + \frac{r_2}{r} + \frac{r^2}{r_1^2} \right) - \left(1 - \frac{r_1}{r} - \frac{r^2}{r_{10}^2} \right) \right\} c^2 dt^2 - \\
& - \left\{ \left(1 + \frac{r_{10}}{r} + \frac{r^2}{r_9^2} \right) - \left(1 - \frac{r_9}{r} - \frac{r^2}{r_8^2} \right) + \left(1 + \frac{r_8}{r} + \frac{r^2}{r_7^2} \right) - \left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2} \right) + \left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2} \right) - \right. \\
& \left. - \left(1 - \frac{r_5}{r} - \frac{r^2}{r_4^2} \right) + \left(1 + \frac{r_4}{r} + \frac{r^2}{r_3^2} \right) - \left(1 - \frac{r_3}{r} - \frac{r^2}{r_2^2} \right) + \left(1 + \frac{r_2}{r} + \frac{r^2}{r_1^2} \right) - \left(1 - \frac{r_1}{r} - \frac{r^2}{r_{10}^2} \right) \right\}^{-1} dr^2 - \\
& - r^2 (d\theta^2 + \sin^2 \theta d\phi^2),
\end{aligned} \tag{67}$$

$$ds_5^{(-)2} = c^2 dt^2 - dr^2 - r^2(d\theta^2 + \sin^2 \theta d\phi^2). \quad (68)$$

Similarly, substitution of series (59) into metrics (49) – (53) leads to the following five similar metrics, but with the opposite signature $(-+++)$:


$$ds_5^{(+)^2} = -c^2 dt^2 + dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2). \quad (69)$$

$$\begin{aligned}
ds_4^{(+2)} = & - \left\{ \left(1 + \frac{r_{10}}{r} + \frac{r^2}{r_9^2}\right) - \left(1 - \frac{r_9}{r} - \frac{r^2}{r_8^2}\right) + \left(1 + \frac{r_8}{r} + \frac{r^2}{r_7^2}\right) - \left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) + \left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) - \right. \\
& \left. - \left(1 - \frac{r_5}{r} - \frac{r^2}{r_4^2}\right) + \left(1 + \frac{r_4}{r} + \frac{r^2}{r_3^2}\right) - \left(1 - \frac{r_3}{r} - \frac{r^2}{r_2^2}\right) + \left(1 + \frac{r_2}{r} + \frac{r^2}{r_1^2}\right) - \left(1 - \frac{r_1}{r} - \frac{r^2}{r_{10}^2}\right) \right\} c^2 dt^2 + \\
& + \left\{ \left(1 + \frac{r_{10}}{r} + \frac{r^2}{r_9^2}\right) - \left(1 - \frac{r_9}{r} - \frac{r^2}{r_8^2}\right) + \left(1 + \frac{r_8}{r} + \frac{r^2}{r_7^2}\right) - \left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) + \left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) - \right. \\
& \left. - \left(1 - \frac{r_5}{r} - \frac{r^2}{r_4^2}\right) + \left(1 + \frac{r_4}{r} + \frac{r^2}{r_3^2}\right) - \left(1 - \frac{r_3}{r} - \frac{r^2}{r_2^2}\right) + \left(1 + \frac{r_2}{r} + \frac{r^2}{r_1^2}\right) - \left(1 - \frac{r_1}{r} - \frac{r^2}{r_{10}^2}\right) \right\}^{-1} dr^2 + \\
& + r^2(d\theta^2 + \sin^2 \theta d\phi^2),
\end{aligned} \tag{70}$$

$$\begin{aligned}
ds_3^{(+2)} = & - \left\{ \left(1 - \frac{r_{10}}{r} - \frac{r^2}{r_9^2}\right) - \left(1 + \frac{r_9}{r} + \frac{r^2}{r_8^2}\right) + \left(1 - \frac{r_8}{r} - \frac{r^2}{r_7^2}\right) - \left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) + \left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) - \right. \\
& \left. - \left(1 + \frac{r_5}{r} + \frac{r^2}{r_4^2}\right) + \left(1 - \frac{r_4}{r} - \frac{r^2}{r_3^2}\right) - \left(1 + \frac{r_3}{r} + \frac{r^2}{r_2^2}\right) + \left(1 - \frac{r_2}{r} - \frac{r^2}{r_1^2}\right) - \left(1 + \frac{r_1}{r} + \frac{r^2}{r_{10}^2}\right) \right\} c^2 dt^2 + \\
& + \left\{ \left(1 - \frac{r_{10}}{r} - \frac{r^2}{r_9^2}\right) - \left(1 + \frac{r_9}{r} + \frac{r^2}{r_8^2}\right) + \left(1 - \frac{r_8}{r} - \frac{r^2}{r_7^2}\right) - \left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) + \left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) - \right. \\
& \left. - \left(1 + \frac{r_5}{r} + \frac{r^2}{r_4^2}\right) + \left(1 - \frac{r_4}{r} - \frac{r^2}{r_3^2}\right) - \left(1 + \frac{r_3}{r} + \frac{r^2}{r_2^2}\right) + \left(1 - \frac{r_2}{r} - \frac{r^2}{r_1^2}\right) - \left(1 + \frac{r_1}{r} + \frac{r^2}{r_{10}^2}\right) \right\}^{-1} dr^2 + \\
& + r^2(d\theta^2 + \sin^2 \theta d\phi^2),
\end{aligned} \tag{71}$$

$$ds_2^{(+)^2} = - \left\{ \left(1 + \frac{r_{10}}{r} - \frac{r^2}{r_9^2} \right) - \left(1 - \frac{r_9}{r} + \frac{r^2}{r_8^2} \right) + \left(1 + \frac{r_8}{r} - \frac{r^2}{r_7^2} \right) - \left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2} \right) + \left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2} \right) - \right. \\ \left. - \left(1 - \frac{r_5}{r} + \frac{r^2}{r_4^2} \right) + \left(1 + \frac{r_4}{r} - \frac{r^2}{r_3^2} \right) - \left(1 - \frac{r_3}{r} + \frac{r^2}{r_2^2} \right) + \left(1 + \frac{r_2}{r} - \frac{r^2}{r_1^2} \right) - \left(1 - \frac{r_1}{r} + \frac{r^2}{r_{10}^2} \right) \right\} c^2 dt^2 + \\ + \left\{ \left(1 + \frac{r_{10}}{r} - \frac{r^2}{r_9^2} \right) - \left(1 - \frac{r_9}{r} + \frac{r^2}{r_8^2} \right) + \left(1 + \frac{r_8}{r} - \frac{r^2}{r_7^2} \right) - \left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2} \right) + \left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2} \right) - \right. \\ \left. - \left(1 - \frac{r_5}{r} + \frac{r^2}{r_4^2} \right) + \left(1 + \frac{r_4}{r} - \frac{r^2}{r_3^2} \right) - \left(1 - \frac{r_3}{r} + \frac{r^2}{r_2^2} \right) + \left(1 + \frac{r_2}{r} - \frac{r^2}{r_1^2} \right) - \left(1 - \frac{r_1}{r} + \frac{r^2}{r_{10}^2} \right) \right\}^{-1} dr^2 + \\ + r^2 (d\theta^2 + \sin^2 \theta d\phi^2),$$

$$\begin{aligned}
ds_1^{(+2)} = & - \left\{ \left(1 - \frac{r_{10}}{r} + \frac{r^2}{r_9^2}\right) - \left(1 + \frac{r_9}{r} - \frac{r^2}{r_8^2}\right) + \left(1 - \frac{r_8}{r} + \frac{r^2}{r_7^2}\right) - \left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) + \left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) - \right. \\
& \left. - \left(1 + \frac{r_5}{r} - \frac{r^2}{r_4^2}\right) + \left(1 - \frac{r_4}{r} + \frac{r^2}{r_3^2}\right) - \left(1 + \frac{r_3}{r} - \frac{r^2}{r_2^2}\right) + \left(1 - \frac{r_2}{r} + \frac{r^2}{r_1^2}\right) - \left(1 + \frac{r_1}{r} - \frac{r^2}{r_{10}^2}\right) \right\} c^2 dt^2 + \\
& + \left\{ \left(1 - \frac{r_{10}}{r} + \frac{r^2}{r_9^2}\right) - \left(1 + \frac{r_9}{r} - \frac{r^2}{r_8^2}\right) + \left(1 - \frac{r_8}{r} + \frac{r^2}{r_7^2}\right) - \left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) + \left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) - \right. \\
& \left. - \left(1 + \frac{r_5}{r} - \frac{r^2}{r_4^2}\right) + \left(1 - \frac{r_4}{r} + \frac{r^2}{r_3^2}\right) - \left(1 + \frac{r_3}{r} - \frac{r^2}{r_2^2}\right) + \left(1 - \frac{r_2}{r} + \frac{r^2}{r_1^2}\right) - \left(1 + \frac{r_1}{r} - \frac{r^2}{r_{10}^2}\right) \right\}^{-1} dr^2 + \\
& + r^2(d\theta^2 + \sin^2 \theta d\phi^2).
\end{aligned} \tag{73}$$



Let's try to connect the radii r_j , included in Ex.s (56) and (57), with the characteristic sizes of bodies in the world around us.

Let's assume that in the basis of a fully geometrized cosmological model there are only two geometric constants: R_v is the parametric radius of the universe, and $l_c \approx c\Delta t \approx c \cdot 1 \text{ sec} \approx 2,9 \cdot 10^{10} \text{ cm}$ is the distance that a light beam travels in vacuum in a time interval $\Delta t = 1 \text{ sec}$.

Let's assume that the radii r_j in the metrics (64) – (73) are estimated by the recursive formula consisting of the above two constants

$$r_j \sim R_v^2 / l_{cj}, \tag{74}$$

where $l_{cj} = (2,9 \cdot 10^{10})^j \text{ cm}$ is the distance obtained by raising the number $2,9 \cdot 10^{10}$ to the power j (1, 2, ..., 10) with the dimension centimeter.

If we assume $R_v \approx 10^{25}$ cm, then we obtain the following approximate recurrent formula

$$r_j \sim \frac{R_v^2}{l_{cj}} = \frac{10^{50}}{(2.9 \cdot 10^{10})^j} \text{ cm}, \quad (75)$$

from which follows a hierarchical sequence of radii of ten spheres nested into each other (see Fig. 5, 6):

(76)

- $r_1 \sim 3.4 \cdot 10^{39}$ cm is the radius commensurate with the radius of the mega-universe*;
- $r_2 \sim 1.2 \cdot 10^{29}$ cm is the radius commensurate with the radius of the core of the metagalaxy (i.e., the observable universe);
- $r_3 \sim 4 \cdot 10^{18}$ cm is the radius commensurate with the radius of the galaxy core;
- $r_4 \sim 1.4 \cdot 10^8$ cm is radius commensurate with the radius of the core of a star (planet);
- $r_5 \sim 4.9 \cdot 10^{-3}$ cm is a radius commensurate with the size of a biological cell;
- $r_6 \sim 1.7 \cdot 10^{-13}$ cm radius commensurate with the core of an elementary particle;
- $r_7 \sim 5.8 \cdot 10^{-24}$ cm is radius, commensurate with the size of a proto-quark*;
- $r_8 \sim 2.1 \cdot 10^{-34}$ cm is radius, commensurate with the size of the plankton nucleus*;
- $r_9 \sim 7 \cdot 10^{-45}$ cm is the radius commensurate with the radius of the proto-plankton core*;
- $r_{10} \sim 2.4 \cdot 10^{-55}$ cm is the radius commensurate with the size of the instanton core*.



Fig. 7. Fractal illustration of two levels of a sequence of nested spherical formations




Fig. 5. Hierarchical sequence of nested spherical formations




Fig. 6. Three levels of a sequence of nested spherical formations

The existence of spherical formations marked with an asterisk * has not been confirmed in practice due to the imperfection of modern technology. Therefore, it is proposed to consider the 10-level cosmological model as a speculative forecast of the author intended for the first working hypothesis. This forecast is based on the confessional intuition of the author, connected with the philosophical doctrine of the "Etz Hayim" ("Tree of Life", i.e., "Tree of ten Sefirot").

However, the radii r_2, r_3, r_4, r_5 and r_6 from the sequence (76), obtained using the recurrent formula (75), surprisingly turned out to be commensurate with the characteristic sizes of the cores (or nuclei) of the observed discrete hierarchical sequence of real spherical formations: metagalaxies, galaxies, stars (planets) and biological cells, while the radius r_6 practically coincided with the "classical radius of the electron" $2.8 \cdot 10^{-13}$ cm. Therefore, it is possible that compact formations with characteristic sizes r_1, r_7, r_8, r_9 and r_{10} can also be detected over time.

Metrics (64) – (73) with a hierarchical sequence of radii r_j (76) describe the metric-dynamic state of a sequence of ten nested multilayer spherical formations (see Figures 5 – 7), the proportions of which partially coincide with the sizes of a discrete sequence of cores (or nuclei) of real objects [17]. Therefore, such a ten-layer and ten-level cosmological model looks promising for further development and refinement.

However, this hierarchical model has one circumstance that does not lend itself to logical understanding. The fact is that from metrics (64) – (73) it follows that the 1-st sphere (whose radius is commensurate with the radius of the mega-universe $r_1 \sim 10^{39}$ cm) is located inside the 10-th sphere (with the instanton size $r_{10} \sim 10^{-55}$ cm). To verify this, follow, for example, the sequence of terms:

$$\begin{aligned} & \left(1 + \frac{r_{10}}{r} - \frac{r^2}{r_9^2}\right) - \left(1 - \frac{r_9}{r} + \frac{r^2}{r_8^2}\right) + \left(1 + \frac{r_8}{r} - \frac{r^2}{r_7^2}\right) - \left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) + \left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) - \\ & - \left(1 - \frac{r_5}{r} + \frac{r^2}{r_4^2}\right) + \left(1 + \frac{r_4}{r} - \frac{r^2}{r_3^2}\right) - \left(1 - \frac{r_3}{r} + \frac{r^2}{r_2^2}\right) + \left(1 + \frac{r_2}{r} - \frac{r^2}{r_1^2}\right) - \left(1 - \frac{r_1}{r} + \frac{r^2}{r_{10}^2}\right) \end{aligned} \quad (77a)$$

and pay attention to the last term, which contains r_1 and r_{10} . This means that the sphere with radius r_1 is inside the sphere with radius r_{10} [17], as can be seen from the previous terms of the same expression. A similar situation takes place in all metrics (64) – (73).

Such isolation of the proposed 10-level cosmological model looks very extravagant and requires additional study and reflection. However, the history of science teaches that sometimes what looks impossible is disproved by experiment. For example, 100 years ago, Einstein and many of his contemporaries could not admit the idea that the part of the universe we observe is expanding with acceleration.

Note that today we do not see only the upper and lower "ends" of the proposed 10-layer and 10-level cosmological model, but in the interval from $r_2 \sim 10^{28}$ cm (scale of the observable universe) to $r_6 \sim 10^{-16}$ cm (intranuclear scale), this model can be useful for solving many problems.

On the other hand, the choice of ten levels (i.e. a closed sequence of 10 nested spheres) is based only on ancient religious Sources (such as the "Sefer Yetzirah" by Avraham Avinu ("The Book of Crea-

tion" by the forefather Avraham), and the Kabbalah of Rabbi Moshe Kordovero, who rigidly insist on only 10 Sefirot*.


The author, of course, understands that the Tree of Ten Sephiroth in the post-Lurian Kabbalah is associated with the 10 Spiritual Levels of the Universe and with the 10 Qualities of the Creative BEGINNING. Nevertheless, it is assumed that what we perceive as the reality of the surrounding world is a projection of Spiritual Conduct on the final results of Creation.

At the same time, this is not Modern Platonism, but the search for a speculative support in the development of the first (a priori) working hypothesis about the structure of a multilayer and multilevel cosmological model.

Of course, in scientific research, we cannot rely only on confessional intuition, therefore, the development and refinement of the proposed multilayer and multilevel cosmological model is possible only on the basis of further development of mathematical analysis and accumulation of empirical data.

In this regard, it is interesting to note that 10 metric layers of the cosmological model under consideration is not an intuitive speculation, but a real total number of possible solutions (44) – (53) of the extended Einstein field equations (43).

Note that today we do not see only the upper and lower "ends" of the proposed 10-layer and 10-level cosmological model, but in the interval from $r_2 \sim 10^{28}$ cm (scale of the observable universe) to $r_6 \sim 10^{-16}$ cm (intra-nuclear scale), this model can be useful for solving many problems.



Name of **GOD** is $\aleph_{(אֵלֶּף)=26} \equiv 10+5+6+5=26$ **י-ה-ו-ה**

Arizal taught: $10^2+5^2+6^2+5^2=100+25+36+25=186=40+100+6+40$ **מִקּוֹם (PLACE)**

5 Metric-dynamic models of «electron» and «positron»

For example, let's select from a hierarchical discrete sequence of spheres nested into each other (see Figures 5 and 6) two mutually opposite spherical formations with a radius $r_6 \sim 10^{-13}$ cm, which corresponds to the characteristic size of the "core" of an elementary particle (in particular cores of the «electron» and «positron»).

All other spherical formations from the considered hierarchical model (64) – (73) are arranged similarly. In this work, the names of the particles are quoted, for example, «electron», since the metric-dynamic models of these spherical formations differ in many respects from the model ideas about these formations in modern physics.

In metrics (64) – (68), let's leave for consideration only those terms that contain radii r_6 . As a result, we obtain the following multilayer metric-dynamic model of a "convex" spherical formation, which we will call «electron»:

$$\begin{aligned} &\text{«ELECTRON»} \\ &\text{"Convex" multilayer spherical formation with} \\ &\text{a signature } (+---), \text{ consisting of:} \end{aligned} \tag{78}$$

The outer shell of the «electron»
in the interval $[r_5, r_6]$ (Figure 8)

$$ds_1^{(+---)2} = \left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \tag{79}$$

$$ds_2^{(+---)2} = \left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \tag{80}$$

$$ds_3^{(+---)2} = \left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \tag{81}$$

$$ds_4^{(+---)2} = \left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2); \tag{82}$$

The core of the «electron»
in the interval $[r_5, r_6]$ (Figure 8)

$$ds_1^{(+---)2} = \left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \tag{83}$$

$$ds_2^{(+---)2} = \left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \tag{84}$$

$$ds_3^{(+---)2} = \left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2), \tag{85}$$

$$ds_4^{(+---)2} = \left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right)} - r^2(d\theta^2 + \sin^2 \theta d\phi^2); \tag{86}$$

The shelt of the «electron»
in the interval $[0, \infty]$

$$ds_5^{(+---)2} = c^2 dt^2 - dr^2 - r^2(d\theta^2 + \sin^2 \theta d\phi^2). \tag{87}$$

where $r_5 \sim 10^{-3}$ cm, $r_6 \sim 10^{-13}$ cm, $r_7 \sim 10^{-24}$ cm.




Fig. 8. A model of a spherical formation (in particular, an «electron») with four clearly defined areas: "outer shell", "abyss" (or "rakia"), "core" and inner "nucleolus" (or "particelle")

Similarly, in the metrics (69) – (73) we also leave only those terms that contain the radii r_6 . As a result, we obtain the following multilayer metric-dynamic model of a "concave" spherical formation, which we will call «positron»:

«POSITRON» (88)

"Concave" multilayer spherical formation with
a signature $(-+++)$, consisting of:

The outer shell of the «positron»

in the interval $[r_5, r_6]$ (negative of the Figure 8)

$$ds_1^{(-+++)^2} = -\left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (89)$$

$$ds_2^{(-+++)^2} = -\left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (90)$$

$$ds_3^{(-+++)^2} = -\left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (91)$$

$$ds_4^{(-+++)^2} = -\left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2); \quad (92)$$

The core of the «positron»
in the interval $[r_5, r_6]$ (negative of the Figure 8)

$$ds_1^{(-+++)^2} = -\left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (93)$$

$$ds_2^{(-+++)^2} = -\left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (94)$$

$$ds_3^{(-+++)^2} = -\left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (95)$$

$$ds_4^{(-+++)^2} = -\left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (96)$$

The shelt of the «positron»

in the interval $[0, \infty]$

$$ds_5^{(-+++)^2} = c^2 dt^2 - dr^2 - r^2(d\theta^2 + \sin^2 \theta d\phi^2). \quad (97)$$

The sets of metrics (78) and (88) differ only in their signature. That is, «electron» and «positron» are completely identical, but antipodal copies of each other. If an «electron» is conventionally called a "convex" spherical formation, then a «positron» is exactly the same conventionally "concave" spherical formation.

The Figure 8 shows a metric-dynamic (i.e., fully geometrized) model of a spherical formation with a radius from the hierarchical sequence (76). In a particular case, an "electron" (and/or its exact antipodal copy, a "positron") has (see Figure 8):

- core with radius $r_6 \sim 10^{-13}$ cm;
- inner "nucleolus" with radius $r_7 \sim 10^{-24}$ cm;
- "outer shell", extending from $r_6 \sim 10^{-13}$ cm to $r_5 \sim 10^{-3}$ cm (or up to $r_4 \sim 10^8$ cm, or up to $r_3 \sim 10^{18}$ cm and etc. depending on which spherical formation the given cors of the «electron» is located inside).

In another case, for example, «planets» (or «anti-planets»): the "core" has a radius $r_4 \sim 10^8$ cm; the inner "nucleolus" has a radius $r_5 \sim 10^{-3}$ cm (or $r_6 \sim 10^{-13}$ cm, etc., depending on which spherical formation is located inside the core of the «planet»), and "outer shell" extends from $r_4 \sim 10^8$ cm to $r_3 \sim 10^{18}$ cm (or up to $r_2 \sim 10^{29}$ cm,




Fig. 9. Fractal illustrations of a multilayer "rakia" separating the "core" of a spherical formation (in particular, an «electron») from its "outer shell"

or up to $r_1 \sim 10^{39}$ cm and depending on, inside which sphere is the core of the «planet»). And so on.

The "shelt" (87) or (97) of a spherical formation begins at its center, and ends at infinity. The "shelt" is a kind of memory of the undeformed state of the considered area of space. The "shelt" does not seem to exist in the curved state of this area of space, but without the components of the metric tensor of the "shelt" $g_{ii}^{0(-)}$ or $g_{ii}^{0(+)}$ it is impossible to determine the deformation, relative elongation, speed and direction of movement of each local section of the object under study (see [16, 17]).

The "abyss" (or "rakia") (see Figures 8 and 9) is a spherical boundary between the "core" and the "outer shell" of any spherical vacuum formation. The "rakia" is the most complex spherical area of the object under study, since it contains sub-layers associated with all higher spheres, inside which is the core of the «electron» (see Figures 5 and 9), as well as sub-layers associated with all the spheres that are inside the same core of the «electron».

A detailed study of the sets of metrics (79) – (87) describing the «electron» and (89) – (97) describing the «positron» is given in [17]. In the same place, taking into account all sixteen signatures (37), metric-dynamic models of almost all elementary particles included in the Standard Model are proposed: all types of «quarks», all types of «neutrinos», «mesons», etc., with the exception of Higgs boson (see Appendix 2 and also <http://metraphysics.ru>, Contents, Chapter 2 Extended Einstein equations, sections 2.9 – 2.14).

It is possible that this article to some extent breaks what is written in verses 6 – 10 of the first chapter of the Book of Genesis (Genesis), the first Book of the TORAH:

וַיֹּאמֶר אֱלֹהִים יְהִי רָקִיעַ בְּתוֹךְ הַמַּיִם וַיְהִי מַבְדִּיל בֵּין מַיִם
לַמַּיִם: וַיַּעַשׂ אֱלֹהִים אֶת־הָרָקִיעַ וַיַּבְדֵּל בֵּין הַמַּיִם אֲשֶׁר מִתַּחַת
לָרָקִיעַ וּבֵין הַמַּיִם אֲשֶׁר מֵעַל לָרָקִיעַ וַיְהִי־כֵן: וַיִּקְרָא אֱלֹהִים
לָרָקִיעַ שָׁמַיִם וַיְהִי־עֶרֶב וַיְהִי־בֹקֶר יוֹם שֵׁנִי:

“And GOD said: “Let there be **רָקִיעַ** (rakia - vault, firmament) in the middle of **מַיִם** (maim - waters) and it will be a division between the waters.” And G-D created rakia and separated the water under the **רָקִיעַ** (rakia) from the water above the rakia, and it became so. And GOD called this **רָקִיעַ** (rakia, arch) the sky and there was evening and there was morning - the second day.

(TORAH, Breishit, 1:6 or Bible, Genesis 1:6)

6 Conclusion and discussion

The article proposes to take into account all 10 Kottler solutions (14) – (25) of Einstein's field equations (8) when constructing a multilayer and multilevel cosmological model.

This statement is supported by the absence of a logical explanation: why only one of these metrics is chosen, and all the others are rejected. Private preferences in choosing one of these ten metrics for one or another cosmological model do not look convincing and lead to insurmountable logical and mathematical problems. Mathematics does not tolerate unreasonable ignoring of any of the solutions of any equation.

In this paper, we consider a multilayer cosmological model, on average, balanced with respect to the Einsteinian vacuum (i.e., "emptiness"), since the sum (or averaging) of all ten solutions (14) – (25) leads to complete zeroing (i.e., to "vacuum balance").

It is shown that the averaging of metric spaces represented by the quadratic forms $ds_i^{(-)2}$ and $ds_i^{(+)2}$ (14) – (25) is connected with the intertwining (i.e., twisting into bundles or braids) of the corresponding affine spaces, represented by linear forms $ds_i^{(-)}$ and $ds_i^{(+)}$.

Moreover, these affine bundles are described by Clifford algebras with the number of generators equal to the number of intertwined linear forms $ds_i^{(-)}$ and $ds_i^{(+)}$ (see [16,17]).

Ten metrics (14) – (25) correspond to 20 conjugate linear forms tied into affine braids, bundles and knots, and when all 16 signatures (37) are taken into account, conjugate linear forms become $16 \times 10 \times 2 = 320$, which form a complex fabric "affine carpet" (see Figure 10).

The mathematics describing the complex "fabric" of space is presented in detail in the "Algebra of signatures" and in "Stochastic metraphysics" [16,17].

The second innovation proposed in this article is related to the increase to infinity of A_i -terms in Einstein's field equations (1).

The restrictions imposed on the sum of A_i -terms made it possible to propose a 10-layered and 10-level (hierarchical) cosmological model.

This model is a hierarchical sequence of nested spheres (see Figure 5) with the corresponding radii r_j (76), which are obtained using the recurrent formula (75) using only two dimensional parameters: $R_v \approx 10^{25}$ cm is parametric the radius of the universe,



Fig. 10. Fractal illustration of an intricately interwoven "affine carpet"

and $l_c \approx 2,9 \cdot 10^{10}$ cm is the distance that a light beam travels in vacuum in a single time interval $\Delta t = 1 \text{ sec}$.

Wherein, part of the radii r_2 , r_3 , r_4 , r_5 and r_6 from the hierarchy (76) turned out to be commensurate with the characteristic sizes of the cores (or nuclei) of the observed discrete sequence of real spherical formations (i.e., cores): metagalaxies, galaxies, stars (planets), biological cells (bacteria) and elementary particles.

In this regard, it is interesting to note that Einstein is translated from German: "ein" - one, "stein" - stone. When Einstein introduced one Λ -term into his equations, he laid one "Stone" in the foundation of general relativity. But this "Stone" turned out to be the size of the universe.

At the end of the article, from the hierarchical solutions (64) – (73), the metric-dynamic models of the «electron» (78) and «positron» (88) are singled out, which are part of the proposed ten-layer and ten-level cosmological model.

Similarly, from solutions (64) – (73) can be singled out metric-dynamic models a «biological cell» with characteristic radius $r_5 \sim 10^{-3}$ cm, a «star» with characteristic core radius $r_4 \sim 10^8$ cm and a «galaxy» with characteristic core radius $r_3 \sim 10^{18}$ cm, etc.

Mathematical techniques that allow extracting various information about local spherical formations from the set of solutions of Einstein's field equations, including a geometrized description of all known force interactions: electrostatic, electromagnetic, weak and nuclear, are presented in the author's work [17] and on the website <http://metraphysics.ru/>.

The advantages of the multilayer and multilevel cosmological model, proposed in this article, include:

- the complete absence of the need to introduce the concept of matter and its energy-momentum tensor as a source of curvature of 4-dimensional space. Here matter is replaced by an infinite number of spherical objects with 10 types of characteristic sizes r_j (76) [17]. In this case, the right side of the Einstein-Hilbert equations (1) naturally vanishes. At the same time, all problems of general relativity disappear, such as, for example, violations of the law of conservation of energy upon transition to other coordinate systems. In this case, Einstein's field equations (8) and (43) are the general covariant expression of the conservation laws [16,17].

- opens the possibility to explore how spherical objects of different scales (i.e., levels) influence each other through the layers in the "rakia" of their cores (or nuclei). For example, how does a sphere with the scale of the observed universe affect the "rakia" of biological cells, or the "rakia" of the core of «electrons», etc.

The disadvantages of the hierarchical 10-layer and 10-level cosmological model proposed in this article include:

- causal and logical inconsistency of this type of closedness of the universe, in which the largest sphere (with the size of the mega-universe $r_1 \sim 10^{39}$ cm) is inside the smallest sphere (with the size of the instanton $r_{10} \sim 10^{-55}$ cm), see, for example, Ex. (77a);

- unreasonable presence of the anthropic principle element in the recurrent formula (75), since the distance $l_c \approx c \cdot 1 \text{ sec} \approx 2,9 \cdot 10^{10}$ cm, which passes a light beam in vacuum in 1 sec (i.e., approximately during the period of oscillation of the human heart);

- an intuitive (i.e., scientifically unfounded) choice of 10 levels (nested spheres) of the cosmological model, despite the fact that today only five discrete levels out of ten are observed.

- the model does not answer the question why we observe a discrete (i.e., in fact, quantum) scale hierarchy of spherical formations (galaxies, stars, biological cells, elementary particles, etc.). The proposed model only "copies" such a discrete-hierarchical manifestation of reality. Obviously, Einstein's field equations (8) and (43) do not contain the possibility of explaining hierarchical discretization (i.e., scaling quantization). This means that Einstein's field equations are not complete. Perhaps, the Einstein-Cartan equations with torsion, or the equations of the geometry of absolute parallelism in a tetrad representation, have such properties. On the other hand, it can be assumed that hierarchical discreteness is not the result of the properties of a differential equation, but is associated with boundary conditions. This is how Maxwell's equations for a waveguide or resonator lead to a discrete spectrum of electromagnetic waves.

At the same time, the multilayer and multilevel cosmological model proposed here is variable, that is, it can be refined as further analysis and accumulation of empirical data.

Acknowledgements

For assistance, I express my deep appreciation and gratitude to the mathematician David Reid (USA, Israel), Ph.D. Tatyana Levi and IT-specialist Alexander Maslov, rabbi Eliezer Reichman.

Appendix 1

The first Einstein field equation and its solutions

The Einstein-Hilbert equation for empty space (i.e., Einstein vacuum) has the form of equation (1) for $T_{ik} = 0$

$$R_{ik} - \frac{1}{2} R g_{ik} = 0, \quad (1.A1)$$

where $R = g^{ik} R_{ik}$ is the scalar curvature;

$$R_{ik} = \frac{\partial \Gamma_{ik}^l}{\partial x^l} - \frac{\partial \Gamma_{il}^l}{\partial x^k} + \Gamma_{ik}^l \Gamma_{lm}^m - \Gamma_{il}^m \Gamma_{mk}^l \text{ is Ricci tensor;}$$

$$\Gamma_{ik}^\lambda = \frac{1}{2} g^{\lambda\mu} \left(\frac{\partial g_{\mu k}}{\partial x^i} + \frac{\partial g_{i\mu}}{\partial x^k} - \frac{\partial g_{ik}}{\partial x^\mu} \right) \text{ is Christoffel symbols.}$$

Combining Eq.s (1.A1) with the contravariant components of the metric tensor g^{ik} , we obtain

$$g^{ik} \left(R_{ik} - \frac{1}{2} R g_{ik} \right) = R - \frac{n}{2} R = 0, \quad (2.A1)$$

where $n = g^{ik} g_{ik}$ is the number of space dimensions.

For any n -dimensional space (except for $n = 2$), equality (2.A1) can be satisfied only for $R = 0$. Therefore, for $n = 4$, Eq. (1.A1) takes the simplest form

$$R_{ik} = 0. \quad (3.A1)$$

We will call this equation the first Einstein vacuum equation, and it is an expression of the conservation laws, since

$$\nabla_j R_{ik} = \frac{\partial R_{ik}}{\partial x^j} = 0, \quad (3a.A1)$$

$$\text{because} \quad \nabla_j 0 = \frac{\partial 0}{\partial x^j} - \Gamma_{ij}^l 0 - \Gamma_{kj}^l 0 = \frac{\partial 0}{\partial x^j} = 0. \quad (3b.A1)$$

In this case, the solutions of Eq.s (3.A1) describe the metric-dynamic state of stable vacuum formations.

Einstein wrote [19]: "The equation of gravity for empty space is the only rationally substantiated case of field theory that can claim to be rigorous."

Solutions of the Eq.s (3.A1) are considered in many works on modern differential geometry and general relativity. However, none of the publications known to the author discusses the relationship between various solutions of this equations, so we will consider it in sufficient detail.

Solutions to Eq.s (3.A1) are sought in a spherical coordinate system in the form of metrics:

$$ds^{(-)2} = e^\nu c^2 dt^2 - e^\lambda dr^2 - r^2 d\theta^2 - r^2 \sin^2 \theta d\varphi^2 \text{ with signature } (+---), \quad (4.A1)$$

$$ds^{(+)2} = -e^\nu c^2 dt^2 + e^\lambda dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2 \text{ with signature } (-+++), \quad (5.A1)$$

where ν and λ are the required functions of t and r .

As a result of substitution of the covariant and contravariant components of the metric tensor from the metric (4.A1) into Eq. (3.A1) for the stationary (i.e., time-independent) state of the "vacuum", a system of three equations is obtained [20]:

$$\nu = -\lambda, \quad (6.A1)$$

$$-e^\nu (\nu'/r + 1/r^2) + 1/r^2 = 0, \quad (7.A1)$$

$$\nu'' + \nu'^2 + 2\nu'/r = 0. \quad (8.A1)$$

The differential Eq. (7.A1) has three solutions:

$$\nu_1 = \ln(h_1 + h_2/r), \quad \nu_2 = \ln(h_1 - h_2/r), \quad \nu_3 = h_3, \quad (9.A1)$$

where h_1, h_2, h_3 are integration constants.

Eq. (8.A1) also has three solutions:

$$\nu_1 = \ln(1 + r_b b/r), \quad \nu_2 = \ln(1 - r_b b/r), \quad \nu_3 = 0, \quad (10.A1)$$

where r_b is the integration constant (the radius of the spherical volume).

For $h_1 = 1$, $h_2 = r_b$ and $h_3 = 0$, the solutions of Eq.s (7.A1) and (8.A1) coincide.

Substituting three possible solutions (10.A1) into the metric (4.A1) we get three metrics with the same signature (+ − − −):

$$ds_1^{(-)2} = (1 - r_b/r)c^2 dt^2 - (1 - r_b/r)^{-1} dr^2 - r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \quad (11.A1)$$

$$ds_2^{(-)2} = (1 + r_b/r)c^2 dt^2 - (1 + r_b/r)^{-1} dr^2 - r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \quad (12.A1)$$

$$ds_3^{(-)2} = c^2 dt^2 - dr^2 - r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2. \quad (13.A1)$$

Performing similar operations with the components of the metric tensor from the metric (5.A1), we obtain three more metrics that also satisfy Eq.s (3.A1), but with the opposite signature (− + + +):

$$ds_1^{(+2)} = - (1 - r_b/r)c^2 dt^2 + (1 - r_b/r)^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \quad (14.A1)$$

$$ds_2^{(+2)} = - (1 + r_b/r)c^2 dt^2 + (1 + r_b/r)^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \quad (15.A1)$$

$$ds_3^{(+2)} = - c^2 dt^2 + dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2. \quad (16.A1)$$

Irreducible into each other metrics (11.A1) – (16.A1) will be called generalized Schwarzschild metrics.

Metrics (11.A1) – (16.A1) describe the metric-dynamic state of the same vacuum region, therefore it is proposed to consider various options for averaging them, despite the fact that Eq.s (3.A1) is non-linear and, as a rule, in such cases, the sum of his decisions is not his own decision.

If the centers of the metrics (11.A1) – (13.A1) and (14.A1) – (16.A1) are aligned, then it is obvious that their sum is equal to zero

$$ds_1^{(-)2} + ds_2^{(-)2} + ds_3^{(-)2} + ds_1^{(+2)} + ds_2^{(+2)} + ds_3^{(+2)} = 0 \cdot c^2 dt^2 + 0 \cdot dr^2 + 0 \cdot d\theta^2 + 0 \cdot \sin^2 \theta d\phi^2 = 0. \quad (17.A1)$$

$$\text{Received metric} \quad ds^{(0)2} = g_{ij}^{(0)} dx^i dx^j, \quad (18.A1)$$

$$\text{where} \quad g_{ij}^{(0)} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (19.A1)$$

is also a trivial solution of the vacuum Eq.s (3.A1).

Thus, contrary to expectations, the addition of six metrics (11.A1) – (16.A1) led to an additional solution to Eq. (3.A1).

Consider now the arithmetic mean of two metrics (11.A1) and (12.A1)

$$ds_{12}^{(-)2} = 1/2 (ds_a^{(-)2} + ds_b^{(-)2}) = c^2 dt^2 - \frac{r^2}{r^2 - r_b^2} dr^2 - r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2. \quad (20.A1)$$

The distance between two points r_1 and r_2 along the length with the signature (+ − − −) in general relativity is determined by the expression

$$r_2 - r_1 = \int_{r_1}^{r_2} \sqrt{-g_{11}^{(-)}} dr. \quad (21.A1)$$

in the case of substitution $g_{11}^{(-)}$ from the averaged metric (20.A1), we obtain

$$r_2 - r_1 = \int_{r_1}^{r_2} \sqrt{-\left(-\frac{r^2}{r^2 - r_b^2}\right)} dr = \int_{r_1}^{r_2} \frac{r dr}{\sqrt{r^2 - r_b^2}} = \sqrt{r^2 - r_b^2} \Big|_{r_1}^{r_2}. \quad (22.A1)$$

Let's first find the value of the segment between the points $r_1 = r_b$ and $r_2 = \infty$:

$$\sqrt{r^2 - r_b^2} \Big|_{r_b}^{\infty} = -\sqrt{-r_b^2} = -\sqrt{-1}r_b = -ir_b. \quad (23.A1)$$

The length of this segment is equal to the radius of the cavity r_b , and the imaginary nature of this result indicates that there is no "vacuum" in the cavity. Outside this cavity from $r_1 = r_b$ to $r_2 = \infty$ we have

$$r_2 - r_1 = \sqrt{r^2 - r_b^2} \Big|_{r_b}^{\infty} = \sqrt{\infty^2 - r_b^2}. \quad (24.A1)$$

In the absence of vacuum deformation, the distance between the points $r_2 = \infty$ and $r_1 = r_b$ is equal $\infty - r_b$, and in the case under consideration it is equal to (24.A1). The difference between these segments is approximately equal to

$$\sqrt{\infty^2 - r_b^2} - (\infty - r_b) \approx r_b. \quad (25.A1)$$

This result shows that the average vacuum extension on the segment $]r_b, \infty[$ is compressed by the value $\sim r_b$, in all radial directions due to the fact that it is displaced from the cavity with radius (25.A1). This result is similar to an air bubble in a liquid (Figure 2.A.1).

The difference between the initial (non-curved) state of a local area of vacuum and its actual (curved) state is determined by the difference [21]

$$ds^{(-)2} - ds^{0(-)2} = (g_{ij}^{(-)} - g_{ij}^{0(-)}) dx^i dx^j \quad (26.A1)$$

where $g_{ij}^{0(-)}$ are the components of the metric tensor of the non-curved vacuum from the metric (13.A1).

The relative elongation of the vacuum region in this case is [21]

$$l^{(-)} = \frac{ds^{(-)} - ds^{0(-)}}{ds^{0(-)}} = \frac{ds^{(-)}}{ds^{0(-)}} - 1, \quad (27.A1)$$

whence it follows [21]

$$ds^{(-)2} = (1 + l^{(-)})^2 ds^{0(-)2}, \quad (28.A1)$$

and

$$l_i^{(-)} = \sqrt{1 + \frac{g_{ii}^{(-)} - g_{ii}^{0(-)}}{g_{ii}^{0(-)}}} - 1 = \sqrt{\frac{g_{ii}^{(-)}}{g_{ii}^{0(-)}}} - 1. \quad (29.A1)$$



Fig.2.A1. Air bubble in liquid




Fig.3.A1. Graph of the function $l_r^{(-)}$ – the relative elongation of the vacuum in the outer shell surrounding the spherical cavity. The calculation was performed at $r_b = 2$, using the software MathCad 15

The uncarved state of the vacuum section under consideration is given by the metric (13.A1), therefore, substituting the components $g_{ii}^{0(-)}$ and $g_{ii}^{(-)}$, respectively, from (13.A1) and (20.A1) into (29.A1), we obtain the relative elongation of the vacuum in each radial direction in the region from r_b to ∞

$$l_t^{(-)} = 0, \quad l_r^{(-)} = \sqrt{\frac{r^2}{r^2 - r_b^2}} - 1, \quad l_\theta^{(-)} = 0, \quad l_\phi^{(-)} = 0. \quad (30.A1)$$

The graph of the function $l_r^{(-)}$ (30.A1) is shown in Figure 3.A1. For $r = r_b$, this function tends to infinity, and for $r < r_b$ it becomes imaginary.

Here we will not discuss the question: – What is inside the cavity with radius r_b , if the vacuum is displaced from there? When considering the second and third Einstein vacuum equations (8) and (43), this problem will be solved by itself.

Thus, averaging the metrics (11.A1) and (11.A1) leads to a metric-dynamic description of a stable vacuum formation of the "air bubble in liquid" (see Figure 3.A1) type, while these metrics alone do not lead to such results.

Averaging the metrics (14.A1) and (15.A1) allows you to get similar results, but with the opposite signature

$$ds_{12}^{(+2)} = 1/2 \left(ds_1^{(+2)} + ds_2^{(+2)} \right) = -c^2 dt^2 + \frac{r^2}{r^2 - r_b^2} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2. \quad (31.II1)$$

$$l_i^{(+)} = \sqrt{1 + \frac{g_{ii}^{(+)} - g_{ii}^{0(+)}}{g_{ii}^{0(+)}}} - 1 = \sqrt{\frac{g_{ii}^{(+)}}{g_{ii}^{0(+)}}} - 1 = \sqrt{\frac{r^2}{r^2 - r_b^2}} - 1. \quad (32.II1)$$

Appendix 2

The Standard Model of Elementary Particles from the Algebra of signatures

1.A2 Models of x_i^+ -«quark» and x_i^- -«antiquark» in the Algebra of signatures

The fundamentals of "Algebra of signatures" and "Stochastic Metraphysics" (that is, fully geometrized physics from the standpoint of Algebra of signatures) are described in [16,17].

This Appendix contains only the main conclusions of the Algebra of signatures related to the Standard Model of elementary particles (Figure 1A2).

Standard Model of Elementary Particles											
three generations of matter (elementary fermions)						three generations of antimatter (elementary antifermions)			interactions / force carriers (elementary bosons)		
	I	II	III	I	II	III					
mass	$\approx 2.2 \text{ MeV}/c^2$	$\approx 1.28 \text{ GeV}/c^2$	$\approx 173.1 \text{ GeV}/c^2$	$\approx 2.2 \text{ MeV}/c^2$	$\approx 1.28 \text{ GeV}/c^2$	$\approx 173.1 \text{ GeV}/c^2$					
charge	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	$-\frac{2}{3}$	$-\frac{2}{3}$	$-\frac{2}{3}$					
spin	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$					
QUARKS	u up	c charm	t top	\bar{u} antiup	\bar{c} anticharm	\bar{t} antitop	g gluon		H higgs		
	$\approx 4.7 \text{ MeV}/c^2$ $-\frac{1}{3}$ $\frac{1}{2}$ d down	$\approx 96 \text{ MeV}/c^2$ $-\frac{1}{3}$ $\frac{1}{2}$ s strange	$\approx 4.18 \text{ GeV}/c^2$ $-\frac{1}{3}$ $\frac{1}{2}$ b bottom	$\approx 4.7 \text{ MeV}/c^2$ $\frac{1}{3}$ $\frac{1}{2}$ \bar{d} antidown	$\approx 96 \text{ MeV}/c^2$ $\frac{1}{3}$ $\frac{1}{2}$ \bar{s} antistrange	$\approx 4.18 \text{ GeV}/c^2$ $\frac{1}{3}$ $\frac{1}{2}$ \bar{b} antibottom	γ photon				
	$\approx 0.511 \text{ MeV}/c^2$ -1 $\frac{1}{2}$ e electron	$\approx 105.66 \text{ MeV}/c^2$ -1 $\frac{1}{2}$ μ muon	$\approx 1.7768 \text{ GeV}/c^2$ -1 $\frac{1}{2}$ τ tau	$\approx 0.511 \text{ MeV}/c^2$ 1 $\frac{1}{2}$ e^+ positron	$\approx 105.66 \text{ MeV}/c^2$ 1 $\frac{1}{2}$ $\bar{\mu}$ antimuon	$\approx 1.7768 \text{ GeV}/c^2$ 1 $\frac{1}{2}$ $\bar{\tau}$ antitau	Z^0 Z ⁰ boson				
LEPTONS	$< 2.2 \text{ eV}/c^2$ 0 $\frac{1}{2}$ ν_e electron neutrino	$< 0.17 \text{ MeV}/c^2$ 0 $\frac{1}{2}$ ν_μ muon neutrino	$< 18.2 \text{ MeV}/c^2$ 0 $\frac{1}{2}$ ν_τ tau neutrino	$< 2.2 \text{ eV}/c^2$ 0 $\frac{1}{2}$ $\bar{\nu}_e$ electron antineutrino	$< 0.17 \text{ MeV}/c^2$ 0 $\frac{1}{2}$ $\bar{\nu}_\mu$ muon antineutrino	$< 18.2 \text{ MeV}/c^2$ 0 $\frac{1}{2}$ $\bar{\nu}_\tau$ tau antineutrino	W^+ W ⁺ boson		W^- W ⁻ boson		
							W^+ W ⁺ boson		W^- W ⁻ boson		

Fig. 1.A2. Elements of the Standard Model of elementary particles

In the above article, it was conditionally assumed that an «electron» is a spherical convexity in empty space (i.e., the Einstein vacuum), which is described by a set of metrics (79) – (87) with the signature $(+ - - -)$; and a «positron» is the exact opposite copy of an «electron», i.e. spherical concavity in the Einstein vacuum, which is described by a set of metrics (89) – (97) with an inverted signature $(- + + +)$.

The Algebra of Signatures takes into account all 16 signatures (37), and the "building material" of the entire variety of observed objects are «quarks» and «antiquarks»:

$$X_i^- \text{«QUARK» or } X_i^+ \text{«ANTIQUARK»} \quad (1.A2)$$

"Convex-concave" multilayer spherical formation
with one of the following signatures:

$$\begin{aligned} & (+ + + +) \quad (+ + + -) \quad (- + + -) \quad (+ + - +) \\ & (- - - +) \quad (- + + +) \quad (- - + +) \quad (- + - +) \\ & (+ - - +) \quad (+ + - -) \quad (+ - - -) \quad (+ - + +) \\ & (- - + -) \quad (+ - + -) \quad (- + - -) \quad (- - - -), \end{aligned} \quad (2.A2)$$

consisting of:

The outer shell of the x_i^+ «quark» or x_i^- «antiquark»
in the interval $[r_5, r_6]$ (Fig. 8)

$$ds_1^{(\pm)2} = \pm \left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) c^2 dt^2 \pm \frac{dr^2}{\left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right)} \pm r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (3.A2)$$

$$ds_2^{(\pm)2} = \pm \left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) c^2 dt^2 \pm \frac{dr^2}{\left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right)} \pm r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (4.A2)$$

$$ds_3^{(\pm)2} = \pm \left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) c^2 dt^2 \pm \frac{dr^2}{\left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right)} \pm r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (5.A2)$$

$$ds_4^{(\pm)2} = \pm \left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) c^2 dt^2 \pm \frac{dr^2}{\left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right)} \pm r^2 (d\theta^2 + \sin^2 \theta d\phi^2); \quad (6.A2)$$

The core of the x_i^+ -«quark» or x_i^- -«antiquark»

in the interval $[r_6, r_7]$ (Fig. 8)

$$ds_1^{(\pm)2} = \pm \left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) c^2 dt^2 \pm \frac{dr^2}{\left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right)} \pm r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (7.A2)$$

$$ds_2^{(\pm)2} = \pm \left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right)} \pm r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (8.A2)$$

$$ds_3^{(\pm)2} = \pm \left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) c^2 dt^2 \pm \frac{dr^2}{\left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right)} \pm r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (9.A2)$$

$$ds_4^{(\pm)2} = \pm \left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) c^2 dt^2 \pm \frac{dr^2}{\left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right)} \pm r^2(d\theta^2 + \sin^2 \theta d\phi^2); \quad (10.A2)$$

The shell of the x_i^+ -«quark» or x_i^- -«antiquark»

in the interval $[0, \infty]$

$$ds_5^{(\pm)2} = \pm c^2 dt^2 \pm dr^2 \pm r^2(d\theta^2 + \sin^2 \theta d\phi^2). \quad (11.A2)$$

where $r_5 \sim 10^{-3}$ cm, $r_6 \sim 10^{-13}$ cm, $r_7 \sim 10^{-24}$ cm.

Signatures, colors and names of all 16 possible x_i^+ -«quarks» and x_i^- -«antiquarks» presented in Table. 1.A2.

Note that within the framework of the Algebra of signatures, these 16 x_i^+ -«quarks» and x_i^- -«antiquarks» refer not only to elementary particles, but also to any other spherical formations from the hierarchy (76). Only instead of radii r_5, r_6, r_7 in metrics (1.Π2) – (11.Π2) it is necessary to substitute accordingly: for bacteria r_4, r_5, r_6 ; for stars r_3, r_4, r_5 ; for galaxies r_2, r_3, r_4 etc.

Table 1.A2

«Quarks»			«Antiquarks»		
Signature type	10 metrics type (1.A2) - (11.A2) with signature:	x_i^+ -«quarks»	10 metrics type (1.A2) - (11.A2) with signature:	x_i^- -«antiquarks»	Color of «quarks» and «antiquarks»
1–3	(+ – – –)	e_y^+ -«quark» («electron»)	(– + + +)	e_y^- -«antiquark» («positron»)	yellow
3–1	(+ + + –)	d_r^+ -«quark»	(– – – +)	d_r^- -«antiquark»	red
	(+ + – +)	d_g^+ -«quark»	(– – + –)	d_g^- -«antiquark»	green
	(+ – + +)	d_b^+ -«quark»	(– + – –)	d_b^- -«antiquark»	blue
2–2	(+ – – +)	u_r^+ -«quark»	(– + + –)	u_r^- -«antiquark»	red
	(+ – + –)	u_g^+ -«quark»	(– + – +)	u_g^- -«antiquark»	green
	(+ + – –)	u_b^+ -«quark»	(– – + +)	u_b^- -«antiquark»	blue
4	(+ + + +)	i_w^+ -«quark»	(– – – –)	i_w^- -«antiquark»	white

For example, let's represent a u_r^- -«antiquark» in expanded form:

$$\mathbf{u_r^-}\text{-«ANTIQUARK»} \quad (12.A2)$$

"Convex-concave" multilayer spherical formation
with the signatures $(- + + -)$, consisting of:

The outer shell of the u_r^- -«antiquark»

in the interval $[r_5, r_6]$ (Fig. 8)

$$ds_1^{(- + + -)^2} = - \left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2} \right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \quad (13.A2)$$

$$ds_2^{(- + + -)^2} = - \left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2} \right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \quad (14.A2)$$

$$ds_3^{(- + + -)^2} = - \left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2} \right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \quad (15.A2)$$

$$ds_4^{(- + + -)^2} = - \left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2} \right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2; \quad (16.A2)$$

The core of the u_r^- -«antiquark»

in the interval $[r_6, r_7]$ (Fig. 8)

$$ds_1^{(- + + -)^2} = - \left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2} \right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2 \quad (17.A2)$$

$$ds_2^{(- + + -)^2} = - \left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2} \right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \quad (18.A2)$$

$$ds_3^{(- + + -)^2} = - \left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2} \right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \quad (19.A2)$$

$$ds_4^{(- + + -)^2} = - \left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2} \right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2} \right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2; \quad (20.A2)$$

The shelt of the u_r^- -«antiquark»

in the interval $[0, \infty]$

$$ds_5^{(- + + -)^2} = -c^2 dt^2 + dr^2 + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2. \quad (21.A2)$$

where $r_5 \sim 10^{-3}$ cm, $r_6 \sim 10^{-13}$ cm, $r_7 \sim 10^{-24}$ cm.

Let's introduce the notation for 16 types of spherical formations: (22.A2)

10* metrics of the form (12.A2) with signature $(+ - - -)$ – yellow e_y^- -«quark» («electron»);

10 metrics of the form (12.A2) with signature $(- + + +)$ – yellow e_y^+ -«antiquark» («positron»);

10 metrics of the form (12.A2) with signature $(+ + + -)$ – red d_r^+ -«antiquark»;

10 metrics of the form (12.A2) with signature $(+ + - +)$ – green d_g^+ -«antiquark»;

10 metrics of the form (12.A2) with signature $(+ - + +)$ – blue d_b^+ -«antiquark»;

10 metrics of the form (12.A2) with signature $(---+)$ – red d_r^- -«quark»;
 10 metrics of the form (12.A2) with signature $(--+-)$ – green d_g^- -«quark»;
 10 metrics of the form (12.A2) with signature $(-+-)$ – blue d_b^- -«quark»;

 10 metrics of the form (12.A2) with signature $(+--+)$ – red u_r^+ -«antiquark»;
 10 metrics of the form (12.A2) with signature $(+-+)$ – green u_g^+ -«antiquark»;
 10 metrics of the form (12.A2) with signature $(++-)$ – blue u_b^+ -«antiquark»;

 10 metrics of the form (12.A2) with signature $(-++-)$ – red u_r^- -«quark»;
 10 metrics of the form (12.A2) with signature $(-++)$ – green u_g^- -«quark»;
 10 metrics of the form (12.A2) with signature $(-++)$ – blue u_b^- -«quark»;

 10 metrics of the form (12.A2) with signature $(----)$ – white i_w^- -«quark»;
 10 metrics of the form (12.A2) with signature $(++++)$ – white i_w^+ -«antiquark».

*10 metrics (12.A2), because Shelt type (21.A2) refers to both the "core" and the "outer shell" of the corresponding x_i^- -«quark» or x_i^+ -«antiquark» (Figure 8). Thus, 5 metrics describe the "core", and 5 metrics describe the "outer shell" of each x_i^- -«quark» and x_i^+ -«antiquark».

Of the spherical formations (22.A2), only a convex formation are stable – «electron» (e_y^- -«quark») with the signature $(+---)$ and a concave formation is a “positron” (e_y^+ -«antiquark») with signature $(-+++)$, because they consist of solutions of Einstein's field equations (8), which is the stability condition. All other spherical formations from the list (22.A2) are unstable, because do not satisfy the conditions of stability (8). That is, when substituting the components of metric tensors from metrics (1.A2) – (11.A2) with any other signature except $(+---)$ and $(-+++)$ into Eq.s (8), equality will not work.

At the same time, out of 16 x_i^+ -«quark» and x_i^- -«antiquark» from the list (22.A2) it is possible to compose averaged stable spherical formations with signatures $(+---)$ or $(-+++)$. This will be shown in the following paragraphs.

2.A2 Models of «proton» and «antiproton» in the Algebra of signatures

On average, stable spherical formations with signatures $(+---)$ or $(-+++)$ can be composed of two different-colored u -«quarks» (or u -«antiquarks») and one d -«quark» (or d -«antiquark») from the list (22.A2):

$$\begin{array}{lll} d_k^+(+ + + -) & d_3^+(+ + - +) & d_r^+(+ - + +) \\ u_3^-(- + - +) & u_r^-(- - + +) & u_k^-(- + + -) \\ \underline{u_r^-(- - + +)} & \underline{u_k^-(- + + -)} & \underline{u_3^-(- + - +)} \\ p_1^-(- + + +)_+ & p_2^-(- + + +)_+ & p_3^-(- + + +)_+ \end{array} \quad \begin{array}{l} (23.A2) \\ (24.A2) \end{array} \quad \begin{array}{l} (25.A2) \end{array}$$

where p_i^- are three possible states of p_i^- -«proton» ($i = 1, 2, 3$) with signature $(-+++)$

$$\begin{array}{lll} d_k^-(- - - +) & d_3^-(- - + -) & d_r^-(- + - -) \\ u_3^+(+ - + -) & u_r^+(+ + - -) & u_k^+(+ - - +) \\ \underline{u_r^+(+ + - -)} & \underline{u_k^+(+ - - +)} & \underline{u_3^+(+ - + -)} \\ p_1^+(+ - - -)_+ & p_2^+(+ - - -)_+ & p_3^+(+ - - -)_+ \end{array} \quad \begin{array}{l} (26.A2) \\ (27.A2) \end{array} \quad \begin{array}{l} (28.A2) \end{array}$$

where p_i^+ are three possible states of p_i^+ -«antiproton» with signature $(+---)$.

In a more compact form, the states p_i^- -«proton» and p_i^+ -«antiproton» can be represented as

$$p_1^+ = u_3^- u_\Gamma^- d_K^+, \quad p_2^+ = u_K^- u_\Gamma^- d_3^+, \quad p_3^+ = u_3^- u_K^- d_\Gamma^+, \quad (29.A2)$$

$$p_1^- = u_3^- u_\Gamma^- d_K^+, \quad p_2^- = u_K^- u_\Gamma^- d_3^+, \quad p_3^- = u_3^- u_K^- d_\Gamma^+. \quad (30.A2)$$

This type of recording of the states of the «proton» and «antiproton» almost completely coincides with the recording of the states of the proton and antiproton in modern quantum chromodynamics, on which the Standard Model of elementary particles is based.

The difference, however, is that in the Standard Model, protons are made up of quarks and antiprotons are made up of antiquarks, while in the Algebra of signatures p_i^- -«proton» and p_i^+ -«antiproton» are made up of x_i^+ -«quarks» and x_i^- -«antiquarks». Therefore, in the Algebra of signatures, there is no problem associated with the baryon asymmetry of the Universe.

For example, let's imagine a multilayer metric-dynamic model of p_i^- -«proton» (23.A2)

$$\begin{array}{c} d_K^+ (+ + + -) \\ u_3^- (- + - +) \\ u_\Gamma^- (- - + +) \\ \hline p_1^+ (- + + +) + \end{array}$$

in the expanded form:

$$\mathbf{p_1^- - «PROTON»} \quad (31.A2)$$

On average, a "concave" multilayer vacuum formation
with a common signature $(- + + +)$
consisting of:

$$\mathbf{d_r^+ - «quark»}$$

"Convex-concave" multilayer spherical formation
with the signatures $(+ + + -)$, consisting of:

$$\mathbf{The\ outer\ shell\ of\ the\ } d_r^+ \text{-«quark» } (+ + + -) \quad (32.A2)$$

in the interval $[r_5, r_6]$ (Fig.1.A2)

$$\begin{aligned} ds_1^{(++++)^2} &= \left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \\ ds_2^{(++++)^2} &= \left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \\ ds_3^{(++++)^2} &= \left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \\ ds_4^{(++++)^2} &= \left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \end{aligned}$$

$$\mathbf{The\ core\ of\ the\ } d_r^+ \text{-«quark» } (+ + + -) \quad (33.A2)$$

in the interval $[r_6, r_7]$ (Fig.1.A2)

$$\begin{aligned} ds_1^{(++++)^2} &= \left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \\ ds_2^{(++++)^2} &= \left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2, \end{aligned}$$

$$ds_3^{(++++)^2} = \left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2,$$

$$ds_4^{(++++)^2} = \left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right)} + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2,$$

The shelt of the d_r^+ -«quark» (+ + + -)

in the interval $[0, \infty]$

$$ds_5^{(++++)^2} = c^2 dt^2 + dr^2 + r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2;$$
(34.A2)

u_g^- -«antiquark»

"Convex-concave" multilayer spherical formation
with the signatures (- + - +), consisting of:

The outer shell of the u_g^- -«antiquark» (- + - +)

in the interval $[r_5, r_6]$ (Fig.1.A2)

(35.A2)

$$ds_1^{(-++-)^2} = -\left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2}\right)} - r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

$$ds_2^{(-++-)^2} = -\left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2}\right)} - r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

$$ds_3^{(-++-)^2} = -\left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2}\right)} - r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

$$ds_4^{(-++-)^2} = -\left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2}\right)} - r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

The core of the u_g^- -«antiquark» (- + - +)

in the interval $[r_6, r_7]$ (Fig.1.A2)

(36.A2)

$$ds_1^{(-++-)^2} = -\left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2}\right)} - r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

$$ds_2^{(-++-)^2} = -\left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2}\right)} - r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

$$ds_3^{(-++-)^2} = -\left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2}\right)} - r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

$$ds_4^{(-++-)^2} = -\left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right) c^2 dt^2 + \frac{dr^2}{\left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2}\right)} - r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

The shelt of the u_g^- -«antiquark» (- + - +)

in the interval $[0, \infty]$

(37.A2)

$$ds_5^{(-++-)^2} = -c^2 dt^2 + dr^2 - r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2;$$

u_b^- -«antiquark»

"Convex-concave" multilayer spherical formation
with the signatures $(- - + +)$, consisting of:

The outer shell of the u_b^- -«antiquark» $(- - + +)$ (38.A2)
in the interval $[r_5, r_6]$ (Fig.1.A2)

$$\begin{aligned} ds_1^{(- - + +)^2} &= - \left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_6}{r} + \frac{r^2}{r_5^2} \right)} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \\ ds_2^{(- - + +)^2} &= - \left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_6}{r} - \frac{r^2}{r_5^2} \right)} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \\ ds_3^{(- - + +)^2} &= - \left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_6}{r} - \frac{r^2}{r_5^2} \right)} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \\ ds_4^{(- - + +)^2} &= - \left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_6}{r} + \frac{r^2}{r_5^2} \right)} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \end{aligned}$$

The core of the u_b^- -«antiquark» $(- - + +)$ (39.A2)
in the interval $[r_6, r_7]$ (Fig.1.A2)

$$\begin{aligned} ds_1^{(- - + +)^2} &= - \left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_7}{r} + \frac{r^2}{r_6^2} \right)} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \\ ds_2^{(- - + +)^2} &= - \left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_7}{r} - \frac{r^2}{r_6^2} \right)} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \\ ds_3^{(- - + +)^2} &= - \left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 - \frac{r_7}{r} - \frac{r^2}{r_6^2} \right)} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \\ ds_4^{(- - + +)^2} &= - \left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2} \right) c^2 dt^2 - \frac{dr^2}{\left(1 + \frac{r_7}{r} + \frac{r^2}{r_6^2} \right)} + r^2 d\theta^2 + \sin^2 \theta d\phi^2, \end{aligned}$$

The shell of the u_g^- -«antiquark» $(- - + +)$ (40.A2)
in the interval $[0, \infty]$

$$ds_5^{(- - + +)^2} = -c^2 dt^2 - dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

where $r_5 \sim 10^{-3}$ cm, $r_6 \sim 10^{-13}$ cm, $r_7 \sim 10^{-24}$ cm.




Fig. 1.A2. Core of the «proton» consists of 3 practically superimposed cores: the core of one d -«quark» and of two cores u -«antiquarks». The inner nucleoli of these 3 "quarks" are in constant chaotic movement and interweaving with each other.

When averaging homogeneous terms in metrics (32.A2) – (40.A2), is obtained a set of metrics (89) – (97) describing the metric-dynamic state of the «positron». However, it should be expected that the radius of the core of the «protons», consisting of the cores of 1 «quark» and 2 «antiquarks», will be greater than the radius of the core of the «positron», because the inner nucleoli of the three «quarks» are difficult to interact, pushing each other away from the common center $r = 0$ (see Figure 1.A2).

The problem of the confinement of three convex-concave spherical formations: d_r^+ -«quark», u_g^- -«antiquark» and u_b^- -«antiquark» is solved by itself, since each x_i^+ -«quark» or x_i^- -«antiquark» from list (22.A2), except e_y^- and e_y^+ , are unstable deformed states of vacuum.

Separately d_r^+ -«quark», u_g^- -«antiquark» and u_b^- -«antiquark» cannot exist for a long time, because the metrics describing them (32.A2) – (40.A2) are not solutions of Einstein's vacuum equation (8). Only together, they form a stable on average "concave" vacuum formation «proton» (see Figure 1.A2), each averaged layer of which satisfies the stability condition (8).

The centers of the «quarks» d_r^+ , u_b^- and u_g^- should wander so chaotically about the common center $r = 0$ and relative to each other (see Figure 1.A2) so that only on average their centers coincide with the common center of the «proton» cores: $\langle r_r \rangle = r = 0$, $\langle r_g \rangle = r = 0$, $\langle r_b \rangle = r = 0$. Therefore, we are

forced to apply not only the metric-dynamic, but also the statistical description of intranuclear processes, which partly considered in [16,17].

The set of metrics (32.A2) – (40.A2) when using the mathematical techniques given in [16,17], allows to extract information about the set of processes and sub-processes occurring both inside the «proton» core and in its "outer shell".

A serious difference between the Algebra of signatures (AS) and the Standard Model of Elementary Particles (SMEP) is that the AS not only largely coincides with the quantum chromodynamics of SMEP, but also extends to spherical objects of any scale.

For example, if in the metrics (1.A2) – (40.A2) we substitute instead of radii:

$$r_5 \sim 10^{-3} \text{ cm}, \quad r_6 \sim 10^{-13} \text{ cm}, \quad r_7 \sim 10^{-24} \text{ cm},$$

respectively, the radii from the hierarchical sequence (76):

$$r_3 \sim 10^{18} \text{ cm}, \quad r_4 \sim 10^8 \text{ cm}, \quad r_5 \sim 10^{-3} \text{ cm},$$

then we get stellar metric Chromodynamics.

Similarly, if from the same sequence (76) we substitute into metrics (32.A2) – (40.A2), respectively, the radii

$$r_2 \sim 10^{29} \text{ cm}, \quad r_3 \sim 10^{18} \text{ cm}, \quad r_4 \sim 10^8 \text{ cm},$$

then we get galactic metric Chromodynamics, and so on.

3.A2 The «neutron» model in the Algebra of signatures

In modern nuclear physics, it is believed that the neutron consists of two d-quarks with a charge of $(-1/3)e$ and one u-quark with a charge of $(2/3)e$ (where e is the electron charge)

$$n = ddu. \quad (41.A2)$$

As a result of this combination, the neutron turns out to be an electrically neutral particle with a zero total charge $(-1/3)e + (-1/3)e + (2/3)e = 0$.

In Algebra of signatures of a «particle» consisting of three «quarks» («antiquarks») with a zero “electric” environment are not obtained. Since there is not a single additive combination of three of the 16 signatures (37), leading to a zero signature (0 0 0 0), which actually means that all vacuum currents (flows) in the outer shell of such a «particle» are completely mutual compensated [17].

The desired result is achieved in the case of rankings consisting of four signatures. Therefore, an "electrically" neutral «particle» («neutron») can have the following topological (nodal) configurations:

$$(42.A2)$$

$i_w^- (- - - -)$	$i_w^- (- - - -)$	$i_w^- (- - - -)$	$i_w^- (- - - -)$
$d_b^+ (+ - + +)$	$d_g^+ (+ + - +)$	$d_b^+ (+ - + +)$	$u_g^- (- + - +)$
$u_r^- (- + + -)$	$d_r^+ (+ + + -)$	$u_g^- (- + - +)$	$d_b^+ (+ - + +)$
$d_g^+ (+ + - +)$	$u_b^- (- - + +)$	$d_r^+ (+ + + -)$	$d_r^+ (+ + + -)$
$n_1^0 (0 0 0 0)_+$	$n_2^0 (0 0 0 0)_+$	$n_3^0 (0 0 0 0)_+$	$n_4^0 (0 0 0 0)_+$

$$\begin{array}{cccc}
i_w^+ (+ + + +) & i_w^+ (+ + + +) & i_w^+ (+ + + +) & i_w^+ (+ + + +) \\
d_g^- (- + - -) & d_g^- (- - + -) & d_b^- (- + - -) & u_g^+ (+ - + -) \\
u_r^+ (+ - - +) & d_r^- (- - - +) & u_g^+ (+ - + -) & d_b^- (- + - -) \\
d_g^- (- - + -) & u_g^+ (+ + - -) & d_r^- (- - - +) & d_r^- (- - - +) \\
n_5^0 (0 0 0 0)_+ & n_6^0 (0 0 0 0)_+ & n_7^0 (0 0 0 0)_+ & n_8^0 (0 0 0 0)_+
\end{array}$$

where

(43.A2)

i_w^+ – white i_w^+ -«quark», i.e. 10 metrics of the form (1.A2) with signature (+ + + +);

i_w^- –white i_w^+ -«antiquark», i.e. 10 metrics of the form (1.A2) with signature (– – – –),

(i from the word *invisible*). These «quarks» are called white because they are practically invisible inside the core of the «neutron», because in terms of topology, they are "point" and "anti-point" [16,17]. Apparently, therefore, their presence in the core of the neutron was not detected experimentally, and was not taken into account by the Standard Model.


Thus, within the framework of the Algebra of signatures, eight possible states of the «neutron» can be represented as:

$$\begin{array}{llll}
n_1^0 = i_6^- d_r^+ d_3^+ u_k^-, & n_2^0 = i_6^- d_k^+ d_3^+ u_r^-, & n_3^0 = i_6^- d_k^+ d_r^+ u_3^-, & n_4^0 = i_6^- d_k^+ d_r^+ u_3^-, \\
n_5^0 = i_6^+ d_r^- d_3^- u_k^+, & n_6^0 = i_6^+ d_3^- d_k^- u_r^+, & n_7^0 = i_6^+ d_r^- d_k^- u_3^+, & n_8^0 = i_6^+ d_r^- d_k^- u_3^+,
\end{array} \quad (44.A2)$$

which differs from the neutron of the Standard Model (41.A2) by the presence of barely distinguishable i_w^+ -«quark» and i_w^- -«antiquark».

Due to the complex topological (or signature) metamorphoses inside the core of the «neutron», any permutation of the 4-«quarks-antiquarks» (44.A2) can be rearranged so that inside a given vacuum formation a combination will be obtained, consisting, for example, of a «proton» and an «electron»:

(45.A2)



Apparently, this restructuring ("decoupling") a topological node inside the core of a «neutron» and leads to a decay reaction

$$n \rightarrow p^+ + e^- + \nu_e, \quad (46.A2)$$

where ν_e is an electronic «neutrino».

4.A2 The model of the «atom» of hydrogen in the Algebra of signatures

Compared to the neutron, the atom of hydrogen is a much more stable formation.

A hydrogen atom (more precisely deuterium) consists of one proton, one neutron and one electron. Within the framework the Algebra of Signature, it also turns out that the deuterium «atom» consists of a «proton», a «neutron» and an «electron». The ranking (topological) equivalent of the nodal configuration of such a spherical formation is:

$$\begin{array}{lcl}
 \begin{array}{l} \text{«proton»} \\ + \\ \text{«neutron»} \\ + \\ \text{«electron»} \\ = \end{array} & \left\{ \begin{array}{l} (+ \ + \ + \ -) \\ (- \ + \ - \ +) \\ (- \ - \ + \ +) \\ (- \ - \ - \ -) \\ (+ \ - \ + \ +) \\ (- \ + \ + \ -) \\ (+ \ + \ - \ +) \\ (+ \ - \ - \ -) \end{array} \right. & \text{or} \quad \left\{ \begin{array}{l} (+ \ + \ - \ +) \\ (- \ - \ + \ +) \\ (- \ + \ + \ -) \\ (+ \ + \ + \ +) \\ (+ \ - \ + \ -) \\ (- \ + \ - \ -) \\ (- \ - \ - \ +) \\ (+ \ - \ - \ -) \end{array} \right. \text{ or } \dots \\
 & & {}^1H(0 \ 0 \ 0 \ 0)_+
 \end{array} \quad (47.A2)$$

Recall that each signature in these rankings corresponds to 10 metrics of the form (1.A2) with a given signature.

The relationship between the metric extent signature and its topology is shown in §1.11 in [17].

It is possible to make many combinations of signatures similar to (47.A2), which reflects the possibilities of "colored" combinatorics of intranuclear metamorphoses. But the topological configuration of this "knot" always remains the same:

$${}^1H = 3u3die, \quad (48.A2)$$

taking into account the topological properties of metrics with corresponding signatures (see §1.11 in [17]), we find that this "knot" consists of 3 intertwined "tori", 4 "oval surfaces" and one "point".

In a similar way, all known chemical elements of the Mendeleev periodic table of elements can be "designed" ("woven"). In this case, the average sizes of the core of «atoms» r_A must depend on the number of x_i^+ -«quarks» and x_i^- -«antiquarks» A that form these "topological nodes"

$$r_A \approx \frac{1}{2} A^{1/3} r_6 \approx \frac{1}{2} A^{1/3} \cdot 10^{-13} \text{ cm}. \quad (49.A2)$$

The Algebra of signatures extends the approach proposed here to the "construction" of multilayer metric-dynamical models of all «atoms» from the periodic table of elements using Fermi's «quarks» and «antiquarks» (with characteristic sizes of core $r_4 \sim 10^8$ cm), to "construction" of metric-dynamical models of "stars» and «planets» with the help of Newton's «quarks» and «antiquarks» (with characteristic sizes of core $r_4 \sim 10^8$ cm), as well as for the construction of «galaxies» with the help of Galileo's «quarks» and «antiquarks» (with characteristic sizes of core $r_3 \sim 10^{18}$ cm), as well as on the construction of «metagalaxies» with the help of Einstein's «quarks» and «antiquarks» (with characteristic sizes of core $r_3 \sim 10^{29}$ cm), etc.

5.A2 Models of «mesons» and «baryons» in the Algebra of signatures

In quantum chromodynamics, mesons are composed of a quark and an antiquark, and are given by the formula

$$M = q^- q^+ = q_{\alpha}^- q_{\alpha}^+ = \frac{1}{\sqrt{3}} (q_b^- q_b^+ + q_r^- q_r^+ + q_g^- q_g^+) g, \quad (50.A2)$$

where q_{α}^- is the color quark triplet ($\alpha = b, g, r$); q_{α}^+ is the color triplet of the antiquark.

Baryons consist of 3 quarks, and are given by the formula

$$B = \frac{1}{\sqrt{6}} q_{\alpha} q_{\beta} q_{\gamma} \varepsilon_{\alpha\beta\gamma}, \quad (51.A2)$$

where $\varepsilon_{\alpha\beta\gamma}$ is a completely antisymmetric tensor.

Almost exactly the same way «mesons» and «baryons» are composed within the Algebra of Signatures. Consider a specific example: three varieties of π -mesons in the theory of strong interactions (quantum chromodynamics) have the following quark structure:

$$\pi^+ = u^- d^+, \quad \pi^0 = \frac{1}{\sqrt{2}} (u^- u^+ - d^+ d^-), \quad \pi^- = u^+ d^-. \quad (52.A2)$$

In the Algebra of signatures, for example, three states of the meson $\pi^+ = u^- d^+$ is represented as

$$\begin{array}{ccc} d_{\kappa}^+ (+ + + -) & d_3^+ (+ + - +) & d_{\Gamma}^+ (+ - + +) \\ \underline{u_3^- (- + - +)} & \underline{u_{\Gamma}^- (- - + +)} & \underline{u_{\kappa}^- (- + + -)} \\ \pi_1^+ (0 \ 2 + 0 \ 0)_+ & \pi_2^+ (0 \ 0 \ 0 \ 2)_+ & \pi_3^+ (0 \ 0 \ 2 + 0)_+ \end{array} \quad (53.A2)$$

where each signature corresponds to a set of 10 metrics of type (1.A2) with a given signature.

In turn, the quark construction

$$\pi^0 = \frac{1}{\sqrt{2}} (u^- u^+ - d^+ d^-) \quad (54.A2)$$

may have the following signature (topological) analogues: (55.A2)

$$\begin{array}{ccc} u_{\kappa}^+ (+ - - +) & u_3^+ (+ - + -) & u_{\Gamma}^+ (+ + - -) \\ u_3^- (- + - +)_+ & u_{\Gamma}^- (- - + +)_+ & u_{\kappa}^- (- + + -)_+ \\ - & - & - \\ d_{\kappa}^+ (+ + + -) & d_3^+ (+ + - +) & d_{\Gamma}^+ (+ - + +) \\ \underline{d_3^- (- - + -)}_+ & \underline{d_{\Gamma}^- (- + - -)}_+ & \underline{d_{\kappa}^- (- - - +)}_+ \\ \pi_1^0 (0 \ 0 \ 0 \ 0) & \pi_2^0 (0 \ 0 \ 0 \ 0) & \pi_3^0 (0 \ 0 \ 0 \ 0) \end{array}$$

Similarly, within the framework of the Algebra of signatures, all known mesons and baryons of the Standard Model can be "constructed".

The construction of the Algebra of signatures (AS) differs from the constructions of the Standard Model of elementary particles, only by the presence of additional i_w^+ -«quark» and i_w^- -«antiquark», as well as by the fact that most of the studied AS of multilayer spherical formations consist of intertwining x_i^+ -«quarks» and x_i^- -«antiquarks», so there is no problem of baryon asymmetry in AS.

6.A2 Models of «bosons» in the Algebra of signatures

«Bosons» are various types of wave (more precisely, helical harmonic) perturbations in vacuum (see §§1.1 – 1.9 in [17]). In this section, we present the main mathematical models of the Algebra of signatures for these wave disturbances.

a) "Photon" and "antiphoton"

Spiral harmonic perturbation [17]

$$\cos\{(2\pi/\lambda)(ct-x-y-z)\} + i \sin\{(2\pi/\lambda)(ct-x-y-z)\} = \exp\{i(2\pi/\lambda)(ct-x-y-z)\} = \exp\{i(\omega t - \mathbf{k} \cdot \mathbf{r})\}. \quad (56.A2)$$

we will conventionally call it a «photon» with signature $\{+ - - -\}$.

Then a spiral harmonic perturbation propagating in the opposite direction, (57.A2)

$$\cos\{(2\pi/\lambda)(-ct+x+y+z)\} + i \sin\{(2\pi/\lambda)(-ct+x+y+z)\} = \exp\{i(2\pi/\lambda)(-ct+x+y+z)\} = \exp\{-i(\omega t - \mathbf{k} \cdot \mathbf{r})\}.$$

we will conventionally call it an «antiphoton» with the signature $\{- + + +\}$.

The notion of the signature of an affine space was introduced in §1.8 in [17].

b) W^\pm -«bosons»

The three color states of the W^+ -«boson» are given by the following expressions and their corresponding rankings [17] (58.A2)

$$\begin{aligned} & \exp\{i2\pi/\lambda(-ct-x-y+z)\} \times & \{- - - +\} \\ & \times \exp\{j2\pi/\lambda(ct-x+y-z)\} \times & \{+ - + -\} \\ & \times \exp\{k2\pi/\lambda(ct+x-y-z)\} & \frac{\{+ + - -\}}{\{+ - - -\}_+} \end{aligned}$$

$$\begin{aligned} & \exp\{i2\pi/\lambda(-ct-x+y-z)\} \times & \{- - + -\} \\ & \times \exp\{j2\pi/\lambda(ct+x-y-z)\} \times & \{+ + - -\} \\ & \times \exp\{k2\pi/\lambda(ct-x-y+z)\} & \frac{\{+ - - +\}}{\{+ - - -\}_+} \end{aligned}$$

$$\begin{aligned} & \exp\{i2\pi/\lambda(-ct+x-y-z)\} \times & \{- + - -\} \\ & \times \exp\{j2\pi/\lambda(ct-x-y+z)\} \times & \{+ - - +\} \\ & \times \exp\{k2\pi/\lambda(ct-x+y-z)\} & \frac{\{+ - + -\}}{\{+ - - -\}_+} \end{aligned}$$

The three color states of the W^- -«boson»:

(59.A2)

$$\begin{aligned} & \exp\{i2\pi/\lambda(ct+x+y-z)\} \times & \{+ + + -\} \\ & \times \exp\{j2\pi/\lambda(-ct+x-y+z)\} \times & \{- + - +\} \\ & \times \exp\{k2\pi/\lambda(-ct-x+y+z)\} & \frac{\{- - + +\}}{\{- + + +\}_+} \end{aligned}$$

$$\begin{aligned} & \exp\{i2\pi/\lambda(ct+x-y+z)\} \times & \{+ + - +\} \\ & \times \exp\{j2\pi/\lambda(-ct-x+y+z)\} \times & \{- - + +\} \\ & \times \exp\{k2\pi/\lambda(-ct+x+y-z)\} & \frac{\{- + + -\}}{\{- + + +\}_+} \end{aligned}$$

$$\begin{aligned}
& \exp \{i 2\pi/\lambda (-ct - x + y + z)\} \times \{+ - + +\} \\
& \times \exp \{j 2\pi/\lambda (-ct + x + y - z)\} \times \{- + + -\} \\
& \times \exp \{k 2\pi/\lambda (-ct + x - y + z)\} \times \frac{\{- + - +\}}{\{- + + +\}}_+,
\end{aligned}$$

where i, j, k are imaginary units, form the anticommutative algebra

$$i^2 = j^2 = k^2 = ijk = -1 \quad \text{и} \quad ij + ji = 0. \quad (60.A2)$$

c) Z^0 -«bosons»

The six color states of the Z^0 -«boson» are given by the following expressions and the corresponding rankings [17] (61.A2)

$$\begin{aligned}
& \exp \{2\pi/\lambda (-ct - x - y - z)\} \times \{- - - -\} \\
& \times \exp \{i 2\pi/\lambda (-ct - x + y + z)\} \times \{+ - + +\} \\
& \times \exp \{j 2\pi/\lambda (-ct + x + y - z)\} \times \{- + + -\} \\
& \times \exp \{k 2\pi/\lambda (-ct + x - y + z)\} \times \frac{\{+ + - +\}}{\{0 \ 0 \ 0 \ 0\}}_+
\end{aligned}$$

$$\begin{aligned}
& \exp \{2\pi/\lambda (-ct - x - y - z)\} \times \{- - - -\} \\
& \times \exp \{i 2\pi/\lambda (-ct + x - y + z)\} \times \{+ + - +\} \\
& \times \exp \{j 2\pi/\lambda (-ct + x + y - z)\} \times \{+ + + -\} \\
& \times \exp \{k 2\pi/\lambda (-ct - x + y + z)\} \times \frac{\{- - + +\}}{\{0 \ 0 \ 0 \ 0\}}_+
\end{aligned}$$

$$\begin{aligned}
& \exp \{2\pi/\lambda (-ct - x - y - z)\} \times \{- - - -\} \\
& \times \exp \{i 2\pi/\lambda (-ct - x + y + z)\} \times \{+ - + +\} \\
& \times \exp \{j 2\pi/\lambda (-ct + x - y + z)\} \times \{- + - +\} \\
& \times \exp \{k 2\pi/\lambda (-ct + x + y - z)\} \times \frac{\{+ + + -\}}{\{0 \ 0 \ 0 \ 0\}}_+
\end{aligned}$$

$$\begin{aligned}
& \exp \{2\pi/\lambda (-ct + x + y + z)\} \times \{+ + + +\} \\
& \times \exp \{i 2\pi/\lambda (-ct + x - y - z)\} \times \{- + - -\} \\
& \times \exp \{j 2\pi/\lambda (-ct - x - y + z)\} \times \{+ - - +\} \\
& \times \exp \{k 2\pi/\lambda (-ct - x + y - z)\} \times \frac{\{- - + -\}}{\{0 \ 0 \ 0 \ 0\}}_+
\end{aligned}$$

$$\begin{aligned}
& \exp \{2\pi/\lambda (-ct + x + y + z)\} \times \{+ + + +\} \\
& \times \exp \{i 2\pi/\lambda (-ct - x + y - z)\} \times \{- - + -\} \\
& \times \exp \{j 2\pi/\lambda (-ct - x - y + z)\} \times \{- - - +\} \\
& \times \exp \{k 2\pi/\lambda (-ct + x - y - z)\} \times \frac{\{+ + - -\}}{\{0 \ 0 \ 0 \ 0\}}_+
\end{aligned}$$

$$\begin{aligned}
& \exp \{2\pi/\lambda (-ct + x + y + z)\} \times \{+ + + +\} \\
& \times \exp \{i 2\pi/\lambda (-ct + x - y - z)\} \times \{- + - -\} \\
& \times \exp \{j 2\pi/\lambda (-ct - x + y - z)\} \times \{+ - + -\} \\
& \times \exp \{k 2\pi/\lambda (-ct - x - y + z)\} \times \frac{\{- - - +\}}{\{0 \ 0 \ 0 \ 0\}}_+
\end{aligned}$$



d) «Graviton» (or «landscapeton»)

In the Algebra of signatures, there is another «boson», which is called «graviton» (or «landscape-ton») [17] (62.A2)

$$\begin{array}{ll}
 \exp \{ \zeta_1 2\pi/\lambda (ct + x + y + z) \} & \{ + \ + \ + \ + \} \\
 \times \exp \{ \zeta_3 2\pi/\lambda (ct - x - y + z) \} \times & \{ - \ - \ - \ + \} \\
 \times \exp \{ \zeta_4 2\pi/\lambda (-ct - x + y - z) \} \times & \{ + \ - \ - \ + \} \\
 \times \exp \{ \zeta_5 2\pi/\lambda (ct + x - y - z) \} \times & \{ - \ - \ + \ - \} \\
 \times \exp \{ \zeta_6 2\pi/\lambda (-ct + x - y - z) \} \times & \{ + \ + \ - \ - \} \\
 \times \exp \{ \zeta_7 2\pi/\lambda (ct - x + y - z) \} \times & \{ - \ + \ - \ - \} \\
 \times \exp \{ \zeta_8 2\pi/\lambda (-ct + x + y + z) \} \times & \{ + \ - \ + \ - \} \\
 \times \exp \{ \zeta_1 2\pi/\lambda (-ct - x - y - z) \} \times & \{ - \ + \ + \ + \} \\
 \times \exp \{ \zeta_2 2\pi/\lambda (ct + x + y - z) \} \times & \{ - \ - \ - \ - \} \\
 \times \exp \{ \zeta_3 2\pi/\lambda (-ct + x + y - z) \} \times & \{ + \ + \ + \ - \} \\
 \times \exp \{ \zeta_4 2\pi/\lambda (ct + x - y + z) \} \times & \{ - \ + \ + \ - \} \\
 \times \exp \{ \zeta_5 2\pi/\lambda (-ct - x + y + z) \} \times & \{ + \ + \ - \ + \} \\
 \times \exp \{ \zeta_6 2\pi/\lambda (ct - x + y + z) \} \times & \{ - \ - \ + \ + \} \\
 \times \exp \{ \zeta_7 2\pi/\lambda (-ct + x - y + z) \} \times & \{ + \ - \ + \ + \} \\
 \times \exp \{ \zeta_8 2\pi/\lambda (ct - x - y - z) \} & \{ - \ + \ - \ + \} \\
 & \underline{\{ + \ - \ - \ - \}} \\
 & \{ 0 \ 0 \ 0 \ 0 \}_+
 \end{array}$$

where the objects ζ_m satisfy the anticommutative relations of the Clifford algebra

$$\zeta_m \zeta_k + \zeta_k \zeta_m = 0 \quad \text{or} \quad m \neq k, \quad \zeta_m \zeta_m = 1, \quad \text{or} \quad \zeta_m \zeta_k + \zeta_k \zeta_m = 2\delta_{km}, \quad (63.A2)$$

where δ_{km} is the Kronecker symbol ($\delta_{km} = 0$ for $m \neq k$ and $\delta_{km} = 1$ for $m = k$). One of the possibilities for determining the objects ζ_m and the Kronecker symbol δ_{km} is presented below:

$$\zeta_5 = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix};$$

$$\zeta_6 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix};$$

$$\zeta_7 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix};$$

$$\zeta_8 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

(64.A2)

7.A2 Conclusions on Annex 2

Metric-dynamic models of all kinds of «neutrinos», «muons» and «antimuons», τ -«leptons», c^+, s^+, b^+, t^+ -«quarks» and c^-, s^-, b^-, t^- -«antiquarks», as well as a geometrized description of the main force interactions: electrostatic, electromagnetic, weak and nuclear are given in chapters 3 – 10 [17].

Thus, taking into account the superposition of stably curved metric spaces with all 16 possible signatures (37)

$$\begin{aligned} & (+ + + +) \quad (+ + + -) \quad (- + + -) \quad (+ + - +) \\ & (- - - +) \quad (- + + +) \quad (- - + +) \quad (- + - +) \\ & (+ - - +) \quad (+ + - -) \quad (+ - - -) \quad (+ - + +) , \\ & (- - + -) \quad (+ - + -) \quad (- + - -) \quad (- - - -) \end{aligned}$$

allow metric-statistical description of almost all elements of the Standard Model of elementary particles, except for the Higgs boson.

In the massless Algebra of signatures (or stochastic metaphysics) proposed here, there is no concept of "mass", so there is no need to introduce the concept of a field that provides a mechanism for spontaneous breaking of electroweak symmetry, and, accordingly, about the quanta of this field - Higgs bosons. However, it is possible that in a fully geometrized theory, metric-dynamic models of vacuum formations with characteristics similar to those of these bosons will arise.

Note that if in the aggregate of metrics of the form (78), (88), (12.A2) and (31.A2) instead of:

$r_5 \sim 10^{-3}$ cm is the characteristic radius of the «biological cell»;

$r_6 \sim 10^{-13}$ cm is the characteristic radius of the «electron» core;

$r_7 \sim 10^{-24}$ cm is the characteristic radius of the «proto-quark» core,

substitute accordingly, for example,

$r_3 \sim 10^{18}$ cm is the radius commensurate with the radius of the «galaxy» core;

$r_4 \sim 10^8$ cm is the radius commensurate with the radius of the core of a «star» («planet»);

$r_5 \sim 10^{-3}$ cm is the radius commensurate with the size of a «biological cell»;

then we get a practically similar multilayer metric-dynamic description of spherical formations on a stellar-planetary scale.

Whereas, if we substitute

$r_2 \sim 10^{29}$ cm is the characteristic radius of the core of the «metagalaxy»;

$r_3 \sim 10^{18}$ cm is the characteristic radius of the core of the «galaxy»;

$r_4 \sim 10^8$ cm – characteristic radius, the core of a "star" or «planet»,

then we get a description of spherical formations of a galactic scale, etc.




Fig. 3.A2. Elements of the Standard Model of elementary particles

Thus, in the opinion of the author, has been obtained a universal metric-dynamic model of a closed and, at the same time, Ricci-flat universe, inhabited by countless spherical formations of various scales,

The probabilistic formalism of the Standard Model also remains valid, since the cores and nucleoli of stable vacuum formations constantly move chaotically under the influence of neighboring stable spherical formations and many other vacuum fluctuations. The study of the chaotic motion of the core of a vacuum formation led to the derivation of the Schrödinger equation (see Chapters 3 and 4 in [17]).

The elements of the Algebra of signatures proposed in this paper are not an alternative theory in relation to general relativity and quantum field theory, but rather their continuation on the path of complete geometrization of physical views.

References

- [1] Einstein, A. (1916) The Foundation of the General Theory of Relativity // *Annalen der Physik*. **354** (7): 769. [Bibcode:1916AnP...354..769E](#). [doi:10.1002/andp.19163540702..](#)
- [2] Einstein, A. (1917) Kosmologische Betrachtungen zur allgemeinen Relativitätstheorie // *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*. Berlin, DE. part 1: pp.142-152. <https://web.archive.org/web/20200329142916/http://echo.mpiwg-berlin.mpg.de/ECHOdocuView?url=/permanent/echo/einstein/sitzungsberichte/S250UZ0K/index.meta>
- [3] Kottler, F. (1918) Über die physikalischen Grundlagen der Einsteinschen Gravitationstheorie// *Annalen der Physik*, Vol. 56, pp. 401-462. [doi:10.1002/andp.19183611402](#).
- [4] Glass, E.N., Krisch, J.P. (2004) Kottler-Lambda-Kerr Spacetime, [arXiv:gr-qc/0405143v1](#) , <https://doi.org/10.48550/arXiv.gr-qc/0405143>.
- [5] [Desai, S.](#) (2022), Galaxy cluster hydrostatic bias in Kottler spacetime// *Physics of the Dark Universe*, Vol. 35, 100928, [arXiv:2111.14105](#)
- [6] [Bambi, C.](#) (2007) Dark Energy and the mass of galaxy clusters. *Phys. Rev. D* **75**, 083003 [arXiv:astro-ph/0703645](#).
- [7] [Nariai, H.](#) (1951). "On a new cosmological solution of Einstein's field equations of gravitation". *Sci. Rep. Tohoku Univ.* **35**: 62.
- [8] [Solanki, R.](#) (2021) Kottler Spacetime in Isotropic Static Coordinates // *Classical and Quantum Gravity*, Vol. 39, No 1, [arXiv:2103.10002](#) , [doi:10.1088/1361-6382/ac3aa0](#).
- [9] [Gibbons, G.W., Warnickand, C.M., Werner, M.C.](#) (2008) "Light-bending in Schwarzschild-de-Sitter: Projective Geometry of the Optical Metric// *Class. Quant. Grav.* **25**, 245009.
- [10] [Cohn, J.D.](#) (1998) Living with Lambda//*Astrophysics and Space Science*, Vol. 259, p. 213–234, <https://doi.org/10.1023/A%3A1001796011627> , [arXiv:astro-ph/9807128v2](#)

- [11] Rindler, W., Ishak, M. (2007) The Contribution of the Cosmological Constant to the Relativistic Bending of Light Revisited// Phys. Rev. D 76, 043006, [arXiv:0709.2948](#).
- [12] Sereno, M. (2007) On the influence of the cosmological constant on gravitational lensing in small systems// Phys. Rev. D 77, 043004, [arXiv:0711.1802](#),
<https://doi.org/10.1103/PhysRevD.77.043004>,
- [13] Ishak, M., Rindler, W., Dossett, J., (2010) More on Lensing by a Cosmological Constant // Mon.Not. R. As- tron. Soc. 403, 2152, 0810.4956. [arXiv:0810.4956](#) ,
<https://doi.org/10.1111/j.1365-2966.2010.16261.x>
- [14] Aghanim N. et al. (Planck Collaboration) (2020). Planck 2018 results. VI. Cosmological parameters (англ.) // Astronomy and Astrophysics. 2020. Vol. 641. P. A6. [arXiv:1807.06209](#)
DOI:<https://doi.org/10.1051/0004-6361/201833910>
- [15] Bachelot, A. (2008) The Dirac system on the Anti-de Sitter Universe//*Communications in Mathematical Physics* vol. 283, pp.127–167, [arXiv:0706.1315](#),
<https://doi.org/10.1007/s00220-008-0564-8>
- [16] Batanov-Gaukhman, M. (2021) The Geometrized Vacuum Physics Based Onthe Algebra of Signature. *Preprints*, 2021060658, doi: 10.20944/preprints202106.0658.v1
<https://www.preprints.org/manuscript/202106.0658/v1>.
- [17] Batanov-Gaukhman, M. (2019) Fully geometrized physics from the standpoint of the Algebra of signature// Lambert, ISBN 978-613-9-45278-1, 2019, C. 415.
Available in English: <http://metraphysics.ru/>
- [19] Einstein A.(1966)// Collection of scientific papers. – M.: Nauka,. v.2, p.789.
- [18] Gaukhman, M.Kh. (2007) Algebra of signatures "NAMES" (orange Alsigna)// ed. Rabbi Gavriel Davidov. - M.: LKI, ISBN-5203-027780-x, available on the Internet URL: <http://alsigna.ru/>, section "Orange Alsigna "NAMES"" [in Russian].
- [20] Landau L.D., Lifshitz E.M. (1971) The Classical Theory of Fields / Course of theoretical physics, V. 2 Translated from the Russian by Hamermesh M. Uni-versity of Minnesota – Pergamon Press Ltd. Oxford, New York, Toronto, Sydney, Braunschwei, p. 387.
- [21] Sedov L.I. (1994) Continuum mechanics. T.1. – M.: Nauka, [in Russian].
- [22] Novikov S.P., Taimanov I.A. (2014) Modern geometric structures and fields. - M.: MTSNMO, 2014. - P. 584 [in Russian].